

Simultaneous proof of the dimensional conjecture and of Jacobi's bound

Abstract

F. Ollivier, LIX, UMR CNRS-École polytechnique n° 7161

Avec le soutien du projet ANR LÉDA (Logistique des Équations Différentielles Algébriques).

Introduction

Jacobi's bound, probably formulated by Jacobi around 1840 [2] is an upper bound on the order of a system of n differential equations f_i in n variables x_j , which is expressed as the *tropical determinant* of the order matrix $A := (a_{i,j} := \text{ord}_{x_j} f_i)$, with the convention $\text{ord}_{x_j} f_i := -\infty$ when f_i is free from x and its derivatives :

$$\max_{\sigma \in S_n} \sum_{i=1}^n a_{i,\sigma(i)}.$$

The result is known to hold for *quasi-regular* systems [3], but remains conjectural in the general case. Cohn was the first to relate it to the dimensional conjecture [1], which claims that the differential codimension of a system of r equations is at most r , showing that Jacobi's bound implies the dimensional conjecture.

During my talk at DART III, I proposed a scheme of proof for the bound, using first a proof of the dimensional conjecture, trying to generalize Ritt's proof for codimension 1 [6], which is based on Puiseux series computations. Then, some kind of reduction process was to be used, mostly based of Jacobi's reduction methods. The dimension properties was crucial there to withdraw components defined by two few equations, or equations satisfying relations.

Working to complete this proof, I had trouble with the computation of Puiseux series that required to introduce some change of variables... allowing to get both results at the same time, thanks to the same reduction process. The result we prove is in fact a little more general.

We define here the order of a component of positive dimension s as the order of its intersection with s generic hyperplanes [5]. Working with a system of r equations is not easy. In fact, it is better to consider an *prime algebraic ideal* of codimension r .

THEOREM 1. — *Let f_i , $1 \leq i \leq r$ be a characteristic set of an algebraic ideal $\mathcal{I}$ of codimension r in $\mathcal{F}\{x_1, \dots, x_n\}$, where $\mathcal{F}$ is a differential field of characteristic 0. Let us denote by $S_{r,n}$ the set of injections from $[1, r]$ to $[1, n]$.*

If $\mathcal{P}$ be a prime component of $\{\mathcal{I}\}$, then :

- i) the differential codimension s of $\mathcal{P}$ is at most $r+p$;*
- ii) if the differential codimension of $\mathcal{P}$ is equal to $r+p$, the order of $\mathcal{P}$ is at most the strong Jacobi number*

$$\mathcal{O} := \max_{\sigma \in S_{r,n}} \sum_{i=1}^s a_{i,\sigma(i)}.$$

If the f_i are just arbitrary equations, we easily reduce to the hypotheses of the theorem by considering the prime components of $\sqrt{[f]}$.

1 Main ideas of the proof.

The most concise way of presenting the proof is to replace the recursive reduction process by a *reductio ad absurdum*. Let us assume that i) or ii) is false. There exist counter-examples such that $n-r$ is minimal, and among them counter-examples with minimal Jacobi number. We will try to work out a contradiction.

Let $B := (\lambda_i + a_{i,j})$ be a *minimal canon* [2, 5] for the order matrix, meaning that (λ_i) is the smallest vector of integers such that B has elements maximal in their columns and located in all different lines and columns. We define $\Lambda := \max_i \lambda_i$, $\alpha_i := \Lambda - \lambda_i$ and $\beta_j := \max_i a_{i,j} - \alpha_i$. We say that some ordering $\prec$ on derivatives is a *Jacobi ordering* if $k_1 - \beta_{j_1} < k_2 - \beta_{j_2}$ implies $x_{j_1}^{(k_1)} \prec x_{j_2}^{(k_2)}$.

We may assume that the f_i are ordered by increasing α_i ; let ϖ be the smallest integer such that :

- A) $f_1, \dots, f_{n-\varpi}$ is a characteristic set of a prime differential ideal $\mathcal{Q}$, for a Jacobi ordering ;
- B) $\mathcal{Q} \cap \mathcal{F}[x_j^{(k)} | 1 \leq j \leq n, 0 \leq k \leq \beta_j + \alpha_{n-\varpi}] \subset \mathcal{P}$.

Now, we may assume further that the system f has been chosen with minimal ϖ , among those with minimal $n-r$ and Jacobi number.

Lemma 2. — *Under the above hypotheses, if ϖ is equal to 0, then $\mathcal{Q} \not\subset \mathcal{P}$; if $\varpi > 0$, then $\varpi < n$ and $\mathcal{J} := \mathcal{Q} \cap \mathcal{F}[x_j^{(k)} | 1 \leq j \leq n, 0 \leq k \leq \beta_j + \alpha_{n-\varpi+1}] \not\subset \mathcal{P}$, which is equivalent to saying that condition B) does not stand for $\varpi - 1$.*

PROOF. — As f_1 is a prime polynomial, it must be the char. set of a prime differential ideal, so $\varpi < n$. If the i_0 first equations f_i are such that $\alpha_i = \alpha_1$, then $f_1, \dots, f_{i_0}$ is also a char. set of a differential prime ideal.

Assume that $\varpi > 0$, that condition B) does not stand for $\varpi - 1$ and that $\alpha_{n-\varpi+1} = \dots = \alpha_{i_0}$. Then, some prime component of the radical of $[f_{n-\varpi+1}, \dots, f_{i_0}] + \mathcal{J}$ must be contained in $\mathcal{P}$. For some Jacobi ordering it has a char. set, of which one may extract a char. set of a prime differential ideal, of the form $f_1, \dots, f_{n-\varpi}, g_{n-\varpi+1}, \dots, g_{i_0}$ —we use here the minimality of $n - r$ and of the Jacobi number. Replacing the corresponding f_i by the g_i , we get a new char. set with a smaller value of ϖ , equal to $n - i_0$: a contradiction to the minimality hypothesis.

Assume now that ϖ is equal to 0. If $\mathcal{Q} \subset \mathcal{P}$, the prime component $\mathcal{P}$ would be equal to $\mathcal{Q}$, of which f is a characteristic set for a Jacobi ordering. This would imply that i) the order of $\mathcal{P}$ is $\mathcal{O}$ and ii) its dimension equal to $n - r$. So, $\mathcal{Q} \not\subset \mathcal{P}$. $\square$

We have shown that possible counter-examples are related to *singular* components of the system $\mathcal{I}$.

2 The singular case.

The idea used to achieve the proof in the singular case is to reduce to the regular one by a suitable change of variables. It may be illustrated by the most simple example of equation $f(x) = x'^2 - 4x = 0$. We introduce a change of variables defined by $y = z'$. The new system $x - y^2 = 0, y(y' - 1) = 0$ is equivalent to $f(x) = 0$, but the main and singular components of this system are now respectively associated to the factors $y' - 1 = 0$ and $y = 0$.

As $\mathcal{P}$ is a component that does not contain $\mathcal{Q}$, we may find a minimal n -uple of integer μ_i such that $[f_i^{(\nu)} | 1 \leq i \leq n - \varpi, 0 \leq \nu \leq \mu_i] : H^1$ contains a polynomial that does not belong to $\mathcal{P}$.

We may now assume that the system has been chosen among those with minimal $n - r$, then minimal Jacobi number, then minimal ϖ , in such a way that $\gamma := \max_{i=1}^r \mu_i$ is minimal. Obviously, it must be greater than 0. We may assume that $\mu_i > 0$ for $1 \leq i \leq s$ and that the leading derivatives of f_i is $x_i^{(\alpha_i + \beta_i)}$. We increase the ground field with r arbitrary functions ζ_i : the new ground field is $\mathcal{F}\langle \zeta \rangle$. We introduce new variables $y_i = x'_i + \zeta_i x_i$.

We start with the prime ideal $\mathcal{I} + [y_i - x'_i - \zeta_i x_i]$ for which $f_i, 1 \leq i \leq n - \varpi$ and $y_i - x'_i - \zeta_i x_i, 1 \leq i \leq r$ is a characteristic set.

The first step is to rewrite the equations defining the new variables as $x'_i = y_i - \zeta_i x_i$, to differentiate them $\alpha_i + \beta_i - 1$ times, so that each derivatives $x'_i, \dots, x_i^{(\alpha_i + \beta_i)}$ are expressed as a linear combinations of $y_i, \dots, y_i^{(\alpha_i + \beta_i - 1)}$ and x_i , that may be substituted in the elements of $\mathcal{I}$ to get a new ideal $\mathcal{I}_1$. This does not change the components of the system.

The next step is to eliminate x'_i in $\mathcal{I}'_1$ using $x'_i = y_i - \zeta_i x_i$, which gives $\mathcal{I}_2$ —that is not an ideal! But one gets a new ideal $\mathcal{I}_1 + \mathcal{I}_2$.

The components are again preserved, but the ideal need not be prime. If $\gamma > 1$, then we just keep the “main” component, that is included in $\mathcal{P}$. The values of $n - r$, the Jacobi number and ϖ are preserved, but it is easily seen that the new value of γ is $\gamma - 1$, which contradicts the minimality of γ .

If $\gamma = 1$, then the components of $\sqrt{\mathcal{Q}_1 + \mathcal{Q}_2}$ that are included in $\mathcal{P}$ have the same $n - r$, but strictly smaller Jacobi number, as they correspond to the former singular components, where H did vanish : a final contradiction that completes the proof.

Références

- [1] COHN (Richard M.), « Order and dimension », *Proc. Amer. Math. Soc.* **87** (1983), n° 1, 1–6.
- [2] JACOBI (Carl Gustav Jacob), “Looking for the order of a system of arbitrary ordinary differential equations”, *AAECC* **20**, (1), 7–32, 2009.
- [3] KONDRATIEVA (Marina Vladimirovna), MIKHALEV (Aleksandr Vasil’evich), PANKRATIEV (Evgeniï Vasil’evich), “Jacobi’s bound for independent systems of algebraic partial differential equations”, *AAECC*, **20**, (1), 65–71, 2009.
- [4] OLLIVIER, (François), “Jacobi’s bound and normal forms computations. A historical survey”, *Differential Algebra And Related Topics*, World Scientific Publishing Company, 2010. ISBN 978-981-283-371-6. To appear.
- [5] OLLIVIER (François) and SADIK (Brahim), “La borne de Jacobi pour une diffiété définie par un système quasi régulier” (Jacobi’s bound for a diffiety defined by a quasi-regular system), *Comptes rendus Mathématique*, **345**, 3, 2007, 139–144.
- [6] RITT (Joseph Fels), *Differential Algebra*, Amer. Math. Soc. Colloq. Publ., vol. 33, A.M.S., New-York, 1950.

1. Where H_f is the product of initials and separants of $f_1, \dots, f_{n-\varpi}$.