

Reduction-based Creative Telescoping for P-recursive Sequences via Integral Bases *

Shaoshi Chen^{1,2}, Lixin Du³, Manuel Kauers³, Rong-Hua Wang⁴

¹KLMM, Academy of Mathematics and Systems Science,
Chinese Academy of Sciences, Beijing, 100190, China.

²School of Mathematical Sciences, University of Chinese Academy of Sciences,
Beijing, 100049, China

³Institute for Algebra, Johannes Kepler University, Linz, A4040, Austria

⁴School of Mathematical Sciences, Tiangong University, Tianjin, 300387, China
schen@amss.ac.cn, lixindumath@gmail.com, manuel.kauers@jku.at, wangronghua@tiangong.edu.cn

Dedicated to the memory of Marko Petkovšek.

Abstract

We propose a way to split a given bivariate P-recursive sequence into a summable part and a non-summable part in such a way that the non-summable part is minimal in some sense. This decomposition gives rise to a new reduction-based creative telescoping algorithm based on the concept of integral bases.

Keywords. Abramov-Petkovšek reduction; P-recursive sequences; Symbolic summation.

1 Introduction

With their book $A = B$ [26], Petkovšek, Wilf, and Zeilberger marked a milestone in the development of symbolic summation. At the beginning of the 1990s, with the appearance of creative telescoping and Petkovšek's algorithm, it seemed that the last word on summation in finite terms had been spoken. The machinery described in the book can decide whether a given definite hypergeometric sum can be written as a linear combination of hypergeometric terms. This is not only a remarkable theoretical result, but it also quickly became an indispensable tool that saves people working in all areas of discrete mathematics from the struggle of simplifying hypergeometric sums by hand. However, from today's perspective, it seems that the pioneering work documented in $A = B$ marks rather the beginning than the end of a period of research on algorithms for summation and integration. In fact, the development of symbolic summation has continued in several directions.

One direction of research concerns the generalization of the techniques to more sophisticated types of sums and integrals [2, 18–20, 24, 27–29, 34, etc.]. Thanks to this line of research, we can deal not only with hypergeometric sums, but also with summands and integrands that are defined in terms of systems of linear differential and difference equations (D-finite functions), summands and integrands that are defined in terms of differential or difference fields (elementary functions, $\Pi\Sigma$ expressions), among other classes of functions.

A second direction of research concerns the refinement of the classical algorithms so as to obtain more precise information about a given summation problem. For example, while Gosper's algorithm [21, 26] for indefinite hypergeometric summation can recognize that a given hypergeometric term is not summable,

*S. Chen was partially supported by the NSFC grants (No. 12271511 and No. 11688101), CAS Project for Young Scientists in Basic Research (Grant No. YSBR-034), the Fund of the Youth Innovation Promotion Association (Grant No. Y2022001), CAS, and the National Key Research and Development Project 2020YFA0712300. L. Du was supported by the Austrian FWF grant P31571-N32. M. Kauers was supported by the Austrian FWF grants P31571-N32 and I6130-N. R. Wang was supported by the NSFC grants (No. 12101449 and No. 12271511), and the Natural Science Foundation of Tianjin, China (No. 22JCQNJC00440).

the Abramov-Petkovšek reduction [4] goes one step further and extracts from any given hypergeometric term a maximal summable part. This is in a way a hypergeometric summation analog to the classical Hermite reduction for indefinite integration of rational functions, which writes any given rational function as the sum of an integrable rational function and a proper rational function with a squarefree denominator.

A third direction of research concerns efficiency. Algorithms for creative telescoping produce a so-called telescoper and a so-called certificate. The certificate tends to be much larger than the telescoper and is often not needed. Starting from 2010, algorithms were developed that can construct a telescoper without also constructing the corresponding certificate. This family of algorithms is now known as “reduction-based telescoping”. It was first proposed for rational function integration [7], where it relies on Hermite reduction. It was then generalized to hyperexponential integrands [8], and later in two different ways to D-finite integrands: one version relies on Lagrange’s identity [9, 32], another on integral bases [12, 15, 17]. Also for summation, the case of rational functions was settled first [16]. Then, using Abramov-Petkovšek reduction, it was formulated for the hypergeometric case [14]. Meanwhile, there are also reduction-based telescoping algorithms for P-recursive functions based on Lagrange’s identity [10, 31].

What is now missing to complete the analogy between the summation case and the integration case is a reduction-based algorithm for P-recursive functions using integral bases. The purpose of the present paper is to introduce such an algorithm. The proposed algorithm is a direct analog of our recent algorithm for D-finite functions [12], with the integral bases for the differential case provided in [6, 23] replaced by the integral bases for the shift case provided in [13] (plus a new notion of integrality at infinity, cf. Sect. 2.2). Although some of the details are somewhat technical, it turns out that there is a full analogy between the integration case and the summation case. At the present stage, this is merely a theoretical result. It remains to be seen how implementations of reduction-based algorithms for D-finite functions or P-recursive sequences based on Lagrange’s identity relate to the corresponding algorithms based on integral bases.

2 Integral bases

Let C be a field of characteristic zero and \bar{C} be the algebraic closure of C . Let $C(x)$ be the field of rational functions in x . The shift operator σ on $C(x)$ is the automorphism such that $\sigma(f(x)) = f(x+1)$ for all $f \in C(x)$. Let $C(x)[S]$ be an Ore algebra, where S is the shift operator with respect to x and satisfies the commutation rule $Sf = \sigma(f)S$ for all $f \in C(x)$. The difference operator $S-1$ is denoted by Δ , which satisfies the rule $\Delta f = \sigma(f)\Delta + \Delta(f)$ for all $f \in C(x)$. Let $L = \ell_0 + \ell_1 S + \dots + \ell_r S^r \in C[x][S]$ with polynomial coefficients $\ell_i \in C[x]$ and $\ell_0 \ell_r \neq 0$. Then r is called the *order* of L . We consider the left $C(x)[S]$ -module $A = C(x)[S]/\langle L \rangle$, where $\langle L \rangle = C(x)[S]L$. When there is no ambiguity, an equivalence class $f + \langle L \rangle$ in A is also denoted by f . Every element of A can be uniquely represented by $f = f_0 + f_1 S + \dots + f_{r-1} S^{r-1}$ with $f_i \in C(x)$. An element $f \in A$ is called *summable* in A if there exists $g \in A$ such that $f = \Delta g$.

2.1 Integral elements at finite places

Let us recall the value functions and integral bases for P-recursive sequences introduced in [13]. Let $\bar{C}((q))$ be the field of formal Laurent series in a new indeterminate q . For each $\alpha \in \bar{C}$, the operator $L = \ell_0 + \ell_1 S + \dots + \ell_r S^r$ acts on a sequence $b: \alpha + \mathbb{Z} \rightarrow \bar{C}((q))$ via

$$(L \cdot b)(z) := \ell_0(z+q)b(z) + \dots + \ell_r(z+q)b(z+r) \quad (1)$$

for all $z \in \alpha + \mathbb{Z}$. This gives an action of the algebra $C(x)[S]$ on the space $\bar{C}((q))^{\alpha+\mathbb{Z}}$ of all sequences $b: \alpha + \mathbb{Z} \rightarrow \bar{C}((q))$. Since $\ell_0 \ell_r \neq 0$, the set $\text{Sol}_\alpha(L) := \{b \in \bar{C}((q))^{\alpha+\mathbb{Z}} \mid L \cdot b = 0\}$ is a $\bar{C}((q))$ -vector space of dimension r . It is called the *solution space* of L at α . Since $q \notin \bar{C}$, we have $\ell_0(z+q)\ell_r(z+q) \neq 0$ for every $z \in \alpha + \mathbb{Z}$. So the sequences $b_1, \dots, b_r: \alpha + \mathbb{Z} \rightarrow \bar{C}((q))$ determined by the recurrence $L \cdot b_i = 0$ and the initial values $b_i(\alpha+j) = \delta_{i,j}$ for $i, j = 1, \dots, r$ form a basis of $\text{Sol}_\alpha(L)$, where $\delta_{i,j}$ is the Kronecker symbol.

The *valuation* $\nu_q(a)$ of a nonzero formal Laurent series $a \in \bar{C}((q))$ is the smallest integer $m \in \mathbb{Z}$ such that the coefficient $[q^m]a$ of q^m in a is nonzero. Set $\nu_q(0) = +\infty$. The *value function* $\text{val}_\alpha: A \rightarrow \mathbb{Z} \cup \{\infty\}$ is defined by

$$\text{val}_\alpha(f) := \min_{b \in \text{Sol}_\alpha(L)} \left(\nu_q((f \cdot b)(\alpha)) - \liminf_{n \rightarrow \infty} \nu_q(b(\alpha - n)) \right) \quad (2)$$

for all $f \in A$. The sequence $f \cdot b$ in (2) is defined as $P_f \cdot b$ via the action (1), if $f = P_f + \langle L \rangle$ for some $P_f \in C(x)[S]$. By convention, set $\infty - \infty = \infty$. An element $f \in A$ is called *(locally) integral* at $\alpha \in \bar{C}$ if $\text{val}_\alpha(f) \geq 0$. Let Z be a subset of \bar{C} . An element $f \in A$ is called *(locally) integral* at Z if $\text{val}_\alpha(f) \geq 0$ for all $\alpha \in Z$, i.e., f is locally integral at all $\alpha \in Z$.

For a fixed $\alpha \in \bar{C}$, the set $C(x)_\alpha = \{p/q \mid p, q \in C[x], q(\alpha) \neq 0\}$ forms a subring of $C(x)$. The set of all elements $f \in A$ that are locally integral at some fixed $\alpha \in \bar{C}$ forms a $C(x)_\alpha$ -module, denoted by \mathcal{O}_α . A basis of this module is called a *local integral basis* at α of A . For $Z \subseteq \bar{C}$, a basis of A is called a *local integral basis* at Z if it is a *local integral basis* at all $\alpha \in Z$. If Z is a finite subset of \bar{C} , a local integral basis at Z exists and the algorithm [13] for computing such a basis for P-recursive sequences is similar to that for algebraic functions [33] and D-finite functions [6, 23].

Let $\alpha \in \bar{C}$. For two elements $\alpha_1, \alpha_2 \in \alpha + \mathbb{Z}$, we say $\alpha_1 < \alpha_2$ if $\alpha_1 - \alpha_2 < 0$. Now we recall some relevant properties of integral bases in [13, Section 5.2]. The lemma gives a way of computing val_α by choosing an appropriate basis of the solution space $\text{Sol}_\alpha(L)$.

Lemma 1. *For a fixed $\alpha \in \bar{C}$, let $\zeta \in \alpha + \mathbb{Z}$ be such that $\ell_0 \ell_r$ has no roots in $\{\eta \in \alpha + \mathbb{Z} \mid \eta < \zeta\}$. Let b_1, \dots, b_r be a basis of $\text{Sol}_\alpha(L)$ defined by the initial values $b_j(\zeta + i - 1) = \delta_{i,j}$ for $i, j = 1, \dots, r$. Then for all $f \in A$ and $\eta \in \alpha + \mathbb{Z}$,*

$$\text{val}_\eta(f) = \min_{j=1}^r \nu_q((f \cdot b_j)(\eta)).$$

Proof. See the argument in [13, Section 5.2, item (A)]. ■

The following lemma arises from a discussion in [13, Section 5.2, item (E)].

Lemma 2. *Let $\zeta \in \alpha + \mathbb{Z}$ with $\alpha \in \bar{C}$ be such that $\ell_0 \ell_r$ has no roots in $Z := \{\eta \in \alpha + \mathbb{Z} \mid \eta < \zeta\}$. Then $\{1, S, \dots, S^{r-1}\}$ is a local integral basis of A at $Z \cup \{\zeta\}$.*

Proof. Let η be an arbitrary but fixed point in $Z \cup \{\zeta\}$ and b_1, \dots, b_r be a basis of $\text{Sol}_\alpha(L)$ such that $b_j(\eta + i - 1) = \delta_{i,j}$ for all $i, j = 1, \dots, r$. Since $\ell_0 \ell_r$ has no roots less than η in $\alpha + \mathbb{Z}$, by Lemma 1 we obtain that for all $f \in A$,

$$\text{val}_\eta(f) = \min_{j=1}^r \nu_q((f \cdot b_j)(\eta)). \quad (3)$$

So $\text{val}_\eta(S^{i-1}) = \min_{j=1}^r \nu_q((S^{i-1} \cdot b_j)(\eta)) = \min_{j=1}^r \nu_q(b_j(\eta + i - 1)) = 0$ and hence S^{i-1} is integral at η for all $i = 1, \dots, r$.

Let $f = f_0 + f_1 S + \dots + f_{r-1} S^{r-1} \in A$ with $f_0, \dots, f_{r-1} \in C(x)$. By the construction of the b_j 's, we have $(f \cdot b_j)(\eta) = \sum_{i=1}^r f_{i-1}(\eta + q) b_j(\eta + i - 1) = f_{j-1}(\eta + q)$ for all $j = 1, \dots, r$. It follows that

$$\min_{j=1}^r \nu_q((f \cdot b_j)(\eta)) = \min_{j=1}^r \nu_q(f_{j-1}(\eta + q)). \quad (4)$$

Assume f is integral at η . Then $\text{val}_\eta(f) \geq 0$ by definition. By (3) and (4), we have $\nu_q(f_{j-1}(\eta + q)) \geq 0$ for all $j = 1, \dots, r$, which implies that $f_{j-1}(x)$ has no pole at η . So $f_{j-1} \in C(x)_\eta$ for all $j = 1, \dots, r$. This proves that $\{1, S, \dots, S^{r-1}\}$ is a local integral basis at η . ■

Local integral bases and shift operators are related as follows.

Lemma 3. *Let $\alpha \in \bar{C}$. Then*

- (i) $f \in A$ is locally integral at α if and only if Sf is locally integral at $\alpha - 1$;
- (ii) $\{\omega_1, \dots, \omega_r\}$ is a local integral basis of A at α if and only if $\{S\omega_1, \dots, S\omega_r\}$ is a local integral basis of A at $\alpha - 1$.

Proof. (i) Clear by $\nu_q(((Sf) \cdot u)(\alpha - 1)) = \nu_q((f \cdot u)(\alpha))$ for all $f \in C(x)[S]$ and $u \in \bar{C}((q))^{\alpha + \mathbb{Z}}$.

(ii) “ \Leftarrow ”: Suppose $f = \sum_{i=1}^r f_i(x) \omega_i$ with $f_i(x) \in C(x)$ is integral at α . By the item (i), we get

$$Sf = S \sum_{i=1}^r f_i(x) \omega_i = \sum_{i=1}^r f_i(x+1) S\omega_i$$

is integral at $\alpha - 1$. Since $\{S\omega_1, \dots, S\omega_r\}$ is a local integral basis at $\alpha - 1$, it follows that $f_i(x+1) \in C(x)_{\alpha-1}$. So $f_i(x) \in C(x)_\alpha$ and hence $\{\omega_1, \dots, \omega_r\}$ is a local integral basis at α .

" \Rightarrow ": Suppose $g = \sum_{i=1}^r g_i(x)S\omega_i$ with $g_i(x) \in C(x)$ is integral at $\alpha - 1$. Since $\ell_0\ell_r \neq 0$, both $\{1, S, \dots, S^{r-1}\}$ and $\{S, S^2, \dots, S^r\}$ are $C(x)$ -vector space bases of A . Then the map sending f to Sf is a bijection from A to itself, because $S \sum_{i=1}^r f_i(x)S^{i-1} = \sum_{i=1}^r f_i(x+1)S^i$ for all $f_i(x) \in C(x)$. There exists $f \in A$ such that $g = Sf$. On the other hand, we have $g = S \sum_{i=1}^r g_i(x-1)\omega_i$. So $f = \sum_{i=1}^r g_i(x-1)\omega_i$. By the item (i) and the assumption that $g = Sf$ is integral at $\alpha - 1$, we obtain that f is integral at α . Since $\{\omega_1, \dots, \omega_r\}$ is a local integral basis at α , it follows that $g_i(x-1) \in C(x)_\alpha$. Thus $g_i(x) \in C(x)_{\alpha-1}$. This completes the proof. \blacksquare

Abramov's dispersion function was introduced in rational summation [1, 25]. It has a feature that the dispersion of the denominator of a rational function strictly increases by one after taking the difference. Now we introduce a local version of dispersion functions and investigate their connection with local integral bases.

Definition 4. Let $\alpha \in \bar{C}$ and $p \in C[x]$. The dispersion of p in $\alpha + \mathbb{Z}$ is defined as

$$\text{disp}_\alpha(p) = \max\{k \in \mathbb{N} \mid \exists \xi \in \alpha + \mathbb{Z} \text{ such that } p(\xi) = p(\xi + k) = 0\}.$$

By convention, $\max \emptyset = -\infty$.

Let $W = (\omega_1, \dots, \omega_r)$ be a vector space basis of A over $C(x)$ and $f = \frac{1}{u} \sum_{i=1}^r a_i \omega_i \in A$ with $a_1, \dots, a_r, u \in C[x]$. We write $f = \frac{aW}{u}$, where $a = (a_1, \dots, a_r) \in C[x]^r$. The vector SW is defined as $(S\omega_1, \dots, S\omega_r)$. Throughout this subsection, we assume $\gcd(a_1, \dots, a_r, u) = 1$. Then u is a denominator of f with respect to W .

Proposition 5. Let $f = \frac{aW}{u}$ and $g = \frac{bW}{v}$, where $a, b \in C[x]^r$, $u, v \in C[x]$. Let $\alpha \in \bar{C}$. Suppose $f = \Delta g$ and $\text{disp}_\alpha(v) \geq 0$.

(i) If W is a local integral basis of A at $\alpha + \mathbb{Z}$, then

$$\text{disp}_\alpha(u) \geq \text{disp}_\alpha(v) + 1.$$

(ii) Let $e \in C[x]$ and $M \in C[x]^{r \times r}$ be such that $SW = \frac{1}{e}MW$. If W is a local integral basis of A at $Z := \{\eta \in \alpha + \mathbb{Z} \mid \eta \leq \beta\}$ with $\beta \in \alpha + \mathbb{Z}$ and β is the only possible root of e in $\alpha + \mathbb{Z}$, then

$$\text{disp}_\alpha((x - \beta)u) \geq 1.$$

This means the polynomial u has a root in $\alpha + \mathbb{Z}$ that is distinct from β .

Proof. (i) A direct calculation yields

$$f = \Delta g = Sg - g = \frac{\sigma(b)}{\sigma(v)}SW - \frac{b}{v}W. \quad (5)$$

Since $\text{disp}_\alpha(v) \geq 0$, the polynomial v has a root in $\alpha + \mathbb{Z}$. Let γ be the minimal root of v in $\alpha + \mathbb{Z}$. Since $x - \gamma \mid v$, we have $\sigma(x - \gamma) = x - (\gamma - 1) \mid \sigma(v)$. So $\gamma - 1$ is a root of $\sigma(v)$. Since W is a local integral basis at $\alpha + \mathbb{Z}$, by Lemma 3 we know that SW is also a local integral basis at $\alpha + \mathbb{Z}$. So both W and SW are local integral bases at $\gamma - 1$. Note that $\sigma(v)$ has a root at $\gamma - 1$, but v does not by the minimality of γ . So $Sg = \frac{\sigma(b)}{\sigma(v)}SW$ is not integral at $\gamma - 1$, but $g = \frac{b}{v}W$ is integral at $\gamma - 1$. Thus $f = Sg - g$ is not integral at $\gamma - 1$. This implies that its denominator u has a root at $\gamma - 1$.

Similarly, let β be the maximal root of v in $\alpha + \mathbb{Z}$. Then v has a root at β , but $\sigma(v)$ does not (otherwise $\beta + 1$ would be a root of v). Since both W and SW are local integral bases at β , it follows that g is not integral at β but Sg is integral at β . So $f = Sg - g$ is not integral at β and hence u has a root at β .

By the definition of dispersion, we have

$$\text{disp}_\alpha(u) \geq \beta - (\gamma - 1) = \beta - \gamma + 1 = \text{disp}_\alpha(v) + 1.$$

(ii) If v has a root in $\alpha + \mathbb{Z}$ that is less than or equal to β , let γ be the minimal root of v in $\alpha + \mathbb{Z}$. Then $\gamma \leq \beta$. Since W is a local integral basis at Z , by Lemma 3 we know that SW is a local integral basis at $Z' = \{\eta - 1 \mid \eta \in Z\}$. So both W and SW are local integral bases at $\gamma - 1$. By the same argument as in (i), we obtain that u has a root at $\gamma - 1$, which is distinct from β .

If v has a root in $\alpha + \mathbb{Z}$ that is greater than β , let γ be the maximal root of v in $\alpha + \mathbb{Z}$. Then $\gamma > \beta$ and γ is not a root of $\sigma(v)$. Substituting $SW = \frac{1}{e}MW$ into (5) yields that

$$f = \Delta g = \left(\frac{\sigma(b)}{\sigma(v)} \frac{1}{e}M - \frac{b}{v} \right) W = \frac{a}{u}W.$$

Note that γ is not a root of $\sigma(v)e$ but γ is a root of v . So u has a root at γ , which is distinct from β . \blacksquare

2.2 Integral elements at infinity

In the differential case, a local integral basis at infinity is used to reduce D-finite functions to integrands with higher valuations at infinity [12]. To achieve the same goal in the shift case, we formulate the notion of local integral bases at infinity for P-recursive sequences, using the framework of valued vector spaces, as in [13].

The operator $L = \ell_0 + \ell_1 S + \dots + \ell_r S^r$ with $\ell_0 \ell_r \neq 0$ admits r linearly independent solutions of the form

$$T = x^{-s} \Gamma(x)^{u/v} \phi^x \exp(p(x^{1/v})) a(x^{-1/v}, \log(x)), \quad (6)$$

where $v \in \mathbb{N} \setminus \{0\}$, $u \in \mathbb{Z}$, $s \in \bar{C}$, $\phi \in \bar{C} \setminus \{0\}$, $a \in \bar{C}[[x]][y]$ and $p \in \bar{C}[x]$ with $\deg_x(p) < v$. Such objects are called *generalized series solutions* of L at infinity, see [22, Section 2.4]. Let R be the set of all \bar{C} -linear combinations of series in the form (6). Then R equipped with the natural addition and multiplication forms a ring. Let $\text{Sol}_\infty(L)$ be the set of all series solutions of L in R . It is called the *solution space* of L at infinity. Then $\text{Sol}_\infty(L)$ is a \bar{C} -vector space of dimension r .

We shall use series solutions to define a value function on A at infinity. The valuation of a series at infinity will be defined in terms of the exponent s in (6). Similar to the D-finite case [23], we use a function ι on \bar{C} to choose an order on these exponents.

Definition 6. Let $\iota : \bar{C} \rightarrow \mathbb{R}$ be a function such that for all $s, s_1, s_2 \in \bar{C}$ and $m \in \mathbb{Q}$,

$$(i) \quad \iota(s_1 + s_2) \geq \iota(s_1) + \iota(s_2);$$

$$(ii) \quad \iota(m + s) = \iota(m) + \iota(s);$$

$$(iii) \quad \iota(m) = m.$$

The valuation $\nu_\infty(t)$ of a term $t := x^{-s} \Gamma(x)^{u/v} \phi^x \exp(p(x^{1/v})) \log(x)^\ell$ is defined as $\iota(s)$. The valuation $\nu_\infty(b)$ of a nonzero series $b \in R$ at infinity is the minimum of the valuations of all the terms appearing in b (with nonzero coefficients). Set $\nu_\infty(0) = \infty$. A series $b \in R$ is called *integral at infinity* if $\nu_\infty(b) \geq 0$.

Example 7. For $C \subseteq \overline{\mathbb{C}(t)}$ with t being a new indeterminate, let $[t^0]s \in \mathbb{C}$ be the constant term of $s \in \bar{C}$ when s is expanded as a Puiseux series around a fixed $\alpha \in \mathbb{C} \cup \{\infty\}$. One choice of $\iota : \bar{C} \rightarrow \mathbb{R}$ is $\iota(s) = \Re([t^0]s)$ for all $s \in \bar{C}$, where $\Re(\cdot)$ is the real part of a complex number. Unless otherwise stated, we shall always assume this choice of ι with $\alpha = \infty$ in the examples given. With this convention, $1, x^{-\sqrt{-1}}, x^{-\sqrt{t}}, x^{-\sqrt{t}-1}$ are integral at infinity, but $x^{-\sqrt{-1}+1}, x^{-\sqrt{t}+1}$ are not.

Lemma 8. The function ν_∞ on R satisfies the following properties: for all $a, b \in R$,

$$(i) \quad \nu_\infty(a) = \infty \text{ if and only if } a = 0;$$

$$(ii) \quad \nu_\infty(ab) \geq \nu_\infty(a) + \nu_\infty(b);$$

$$(iii) \quad \nu_\infty(a + b) \geq \min\{\nu_\infty(a), \nu_\infty(b)\}.$$

So the set $\{b \in R \mid b \text{ is integral}\}$ forms a subring of R .

Proof. (i) and (iii) are clear by definition. For (ii), note that $x^{-s_1} x^{-s_2} = x^{-(s_1+s_2)}$ and $\iota(s_1 + s_2) \geq \iota(s_1) + \iota(s_2)$ for all $s_1, s_2 \in \bar{C}$ by the assumption on ι in Definition 6. ■

The shift $\sigma : b(x) \mapsto b(x+1)$ of series is defined through the rules:

$$\begin{aligned}
(x+1)^{-s} &= \sum_{n=0}^{\infty} \binom{-s}{n} x^{-s-n} = x^{-s} - sx^{-s-1} + \dots, \\
\Gamma(x+1)^{u/v} &= x^{u/v} \Gamma(x)^{u/v}, \\
\phi^{x+1} &= \phi \phi^x, \\
\exp(c(x+1)^{k/v}) &= \exp(cx^{k/v}) \exp(c((x+1)^{k/v} - x^{k/v})) \\
&= \exp(cx^{k/v}) \sum_{n=0}^{\infty} \frac{1}{n!} \left(c \sum_{i=1}^{\infty} \binom{k/v}{i} x^{k/v-i} \right)^n \\
&= \exp(cx^{k/v}) \left(1 + \frac{ck}{v} x^{k/v-1} + \dots \right) \quad (c \in \bar{C}, 0 \leq k < v), \\
\log(x+1)^\ell &= \left(\log(x) + \log\left(1 + \frac{1}{x}\right) \right)^\ell = \left(\log(x) - \sum_{n=1}^{\infty} \frac{(-1)^n}{n} x^{-n} \right)^\ell \\
&= \log(x)^\ell + \ell x^{-1} \log(x)^{\ell-1} + \dots,
\end{aligned}$$

where “ \dots ” denotes some terms of higher valuations at infinity.

An operator $P = p_0 + p_1 S + \dots + p_{r-1} S^{r-1}$ in $C(x)[S]$ acts on a series b via

$$P \cdot b = p_0 b + p_1 \sigma(b) + \dots + p_{r-1} \sigma^{r-1}(b).$$

For a solution $b \in \text{Sol}_\infty(L) \subseteq R$ and $f \in A = C(x)[S]/\langle L \rangle$, the action $f \cdot b$ is defined as $P_f \cdot b$ if $f = P_f + \langle L \rangle$ for some $P_f \in C(x)[S]$. Let b_1, \dots, b_r be a basis of $\text{Sol}_\infty(L)$ in the form of (6). The *value function* $\text{val}_\infty : A \rightarrow \mathbb{R} \cup \{\infty\}$ is defined as

$$\text{val}_\infty(f) := \min_{i=1}^r \nu_\infty(f \cdot b_i).$$

Then by Lemma 8, $\text{val}_\infty(f)$ is the minimum valuation of all series $f \cdot b$ at infinity, where b runs through series solutions in $\text{Sol}_\infty(L)$. An element $f \in A$ is called (*locally*) *integral* at infinity if $\text{val}_\infty(f) \geq 0$.

Proposition 9. *The function val_∞ satisfies the following properties: for all $f, g \in A$ and $a \in C(x)$,*

- (i) $\text{val}_\infty(f) = \infty$ if and only if $f = 0$;
- (ii) $\text{val}_\infty(af) = \nu_\infty(a) + \text{val}_\infty(f)$;
- (iii) $\text{val}_\infty(f+g) \geq \min\{\text{val}_\infty(f), \text{val}_\infty(g)\}$;

where $\nu_\infty(a)$ is the valuation of its Laurent series expansion at infinity.

Proof. (i) Let b_1, \dots, b_r be a basis of $\text{Sol}_\infty(L)$ in the form of (6). If $f = 0$, then $f \cdot b_i$ is the zero series for all $i = 1, \dots, r$. So $\text{val}_\infty(f) = \infty$. Conversely, assume that $\text{val}_\infty(f) = \infty$, then by definition $\nu_\infty(f \cdot b_i) = \infty$ for all $i = 1, \dots, r$. So $f \cdot b_i = 0$ for $i = 1, \dots, r$, which implies that f has at least r linearly independent series solutions. On the other hand, every nonzero element of A can be written as $f = f_0 + f_1 S + \dots + f_k S^k$ with $f_i \in C[x]$, $f_k \neq 0$ and $k < r$. If the trailing coefficient f_0 of f is nonzero, the dimension of the solution space of f in R can not exceed its order k , see [22, Section 2.4]. So in this case, f cannot be nonzero.

If $f_0 = 0$, we write $f = Sg$ with $g \in A$ such that the order of g is $k-1$. Then $Sg \cdot b_i = 0$ and we claim that $g \cdot b_i = 0$ for all $i = 1, \dots, r$. Otherwise, suppose $g \cdot b_i \neq 0$ for some $i \in \{1, \dots, r\}$. Then $g \cdot b_i$ is a generalized series involves a term

$$T = x^{-s} \Gamma(x)^{u/v} \phi^x \exp(p(x^{1/v})) \log(x)^\ell,$$

where $v \in \mathbb{N} \setminus \{0\}$, $u \in \mathbb{Z}$, $\ell \in \mathbb{N}$, $s \in \bar{C}$, $\phi \in \bar{C} \setminus \{0\}$ and $p \in \bar{C}[x]$ with $\deg_x(p) < v$. Let T be the dominant term of $g \cdot b_i$, i.e., among all terms with minimal s the one with the largest exponent ℓ . Then

$$S \cdot T = (x+1)^{-s} \Gamma(x+1)^{u/v} \phi^{x+1} \exp(p((x+1)^{1/v})) \log(x+1)^\ell = \phi x^{u/v} T + \dots.$$

Here $\phi \neq 0$ by the assumption $\ell_0 \ell_r \neq 0$. So $\phi x^{u/v} T$ is the dominant term of $Sg \cdot b_i$ and hence $Sg \cdot b_i \neq 0$. This leads to a contradiction. Thus $g \cdot b_i = 0$ for all $i = 1, \dots, r$. Let j with $1 \leq j \leq k$ be the minimal integer such that $f_j \neq 0$. We can write $f = S^j \tilde{g}$ with $\tilde{g} \in A$ such that the order of \tilde{g} is $k - j$ and the trailing coefficient of \tilde{g} is nonzero. We proceed similarly with $g = S^{j-1} \tilde{g}$, and finally obtain that $\tilde{g} \cdot b_i = 0$ for all $i = 1, \dots, r$. So \tilde{g} cannot be nonzero by the same argument as in the first case. Thus f cannot be nonzero.

(ii) Every nonzero $a \in C(x) \subseteq C[[x^{-1}]]$ can be expanded as a Laurent series in the form $a = a_m x^{-m} + a_{m-1} x^{-m-1} + \dots$, where $a_i \in C$, $m \in \mathbb{Z}$ and $a_m \neq 0$. By Definition 6, $\nu_\infty(a) = \iota(m) = m$ is an integer. Note that for any $m \in \mathbb{Z}$ and $s \in \bar{C}$, $x^{-m} x^{-s} = x^{-m-s}$ and $\iota(m+s) = \iota(m) + \iota(s) = m + \iota(s)$. So $\nu_\infty(ab) = \nu_\infty(a) + \nu_\infty(b)$ for all $a \in C(x)$ and $b \in R$. Thus (ii) is true.

(iii) Clear by the item (iii) in Lemma 8. ■

For a rational function p/q with $p, q \in C[x]$, we have $\nu_\infty(p/q) = \deg_x(q) - \deg_x(p)$. The set $C(x)_\infty = \{p/q \mid p, q \in C[x], \deg_x(p) \leq \deg_x(q)\}$ forms a subring of $C(x)$. By Proposition 9, (A, val_∞) is a valued vector space over the valued field $(C(x), \nu_\infty)$, see [13, Definition 2]. So by Proposition 4 in [13], the set of all elements $f \in A$ that are locally integral at infinity forms a $C(x)_\infty$ -module, denoted by \mathcal{O}_∞ . A basis of this module is called a *local integral basis* at infinity of A . Similar to the algebraic and D-finite cases [6, 13, 23, 33], such a basis can be computed by Algorithm 10 in [13]. The termination of this algorithm is guaranteed by the existence of discriminant functions. For a basis $\omega_1, \dots, \omega_r$ of A , the discriminant function Disc is defined as

$$\text{Disc}(\omega_1, \dots, \omega_r) = \lfloor \nu_\infty(\det((\omega_i \cdot b_j)_{i,j=1}^r)) \rfloor,$$

where b_1, \dots, b_r is a basis of $\text{Sol}_\infty(L)$ in the form of (6). It can be checked that the Disc is indeed a discriminant function in [13, Definition 12].

A $C(x)$ -vector space basis $\{\omega_1, \dots, \omega_r\}$ of $A = C(x)[S]/\langle L \rangle$ is called *normal at infinity* if there exist $a_1, \dots, a_r \in C(x)$ such that $\{a_1 \omega_1, \dots, a_r \omega_r\}$ is a local integral basis at infinity. Given a vector space basis, Trager's algorithm [30] can be literally adapted to compute a basis that is normal at infinity and generates the same $C[x]$ -module as the given basis.

For a nonconstant polynomial $g \in C[x]$, we have $\deg_x(\Delta(g)) = \deg_x(g) - 1$. For a rational function $g \in C(x)$, if $\nu_\infty(g) \neq 0$, then $\nu_\infty(\Delta(g)) = \nu_\infty(g) + 1$. So the valuation at infinity of such a rational function increases by exactly one under each difference operation. In the P-recursive case, we shall prove that the valuation at infinity increases by at most one.

The product rule in the shift case says that $\Delta(uv) = \Delta(u)v + \sigma(u)\Delta(v)$ for every $u, v \in R$. For several elements $u_1, \dots, u_n \in R$, it generalizes to

$$\Delta(u_1 \cdots u_n) = \sum_{i=1}^n \sigma(u_1 \cdots u_{i-1}) \Delta(u_i) u_{i+1} \cdots u_n. \quad (7)$$

Proposition 10. *Let $g \in A$. If $\text{val}_\infty(g) \neq 0$, then $\text{val}_\infty(\Delta g) \leq \text{val}_\infty(g) + 1$.*

Proof. Let b_i be a generalized series solution in $\text{Sol}_\infty(L)$ such that $\text{val}_\infty(g) = \nu_\infty(g \cdot b_i)$. There $g \cdot b_i$ involves a term

$$T = x^{-s} \Gamma(x)^{u/v} \phi^x \exp(p(x^{1/v})) \log(x)^\ell,$$

where $v \in \mathbb{N} \setminus \{0\}$, $u \in \mathbb{Z}$, $\ell \in \mathbb{N}$, $s, \phi \in \bar{C} \setminus \{0\}$ and $p \in \bar{C}[x]$ with $\deg_x(p) < v$. For this fixed series b_i , let T be the dominant term of $g \cdot b_i$, i.e., among all terms with minimal r the one with the largest exponent ℓ . Let $k = \deg_x(p)$ and $c = \text{lc}_x(p)$ be the degree and the leading coefficient of p in x respectively (take $k = 0$ if $p(x) = 0$).

Now we use Equation (7) to calculate $\Delta \cdot T$. Note that

$$\begin{aligned} \Delta(x^{-s}) \Gamma(x)^{u/v} \phi^x \exp(p(x^{1/v})) \log(x)^\ell &= -s x^{-1} T + \dots, \\ \sigma(x^{-s}) \Delta(\Gamma(x)^{u/v}) \phi^x \exp(p(x^{1/v})) \log(x)^\ell &= (x^{u/v} - 1) T + \dots, \\ \sigma(x^{-s} \Gamma(x)^{u/v}) \Delta(\phi^x) \exp(p(x^{1/v})) \log(x)^\ell &= (\phi - 1) x^{u/v} T + \dots, \\ \sigma(x^{-s} \Gamma(x)^{u/v} \phi^x) \Delta(\exp(p(x^{1/v}))) \log(x)^\ell &= \frac{\phi c k}{v} x^{u/v+k/v-1} T + \dots, \\ \sigma(x^{-s} \Gamma(x)^{u/v} \phi^x \exp(p(x^{1/v}))) \Delta(\log(x)^\ell) &= \phi \ell x^{u/v-1} \log(x)^{-1} T + \dots. \end{aligned}$$

So by Equation (7), we get

$$\Delta \cdot T = (-sx^{-1} - 1 + \phi x^{u/v} + \frac{\phi ck}{v} x^{u/v+k/v-1})T + \dots$$

Note that $0 \leq \frac{k}{v} < 1$ because $0 \leq k < v$. So $-\frac{u}{v} < -\frac{u}{v} - \frac{k}{v} + 1$. Let T_0 be the dominant term of $(\Delta g) \cdot b_i$. We make the following case distinction.

- (1) If $u > 0$, then $-u/v < \min \{1, 0, -\frac{u}{v} - \frac{k}{v} + 1\}$. We have $T_0 = \phi x^{u/v} T$. Here $\phi \neq 0$ by the assumption $\ell_0 \ell_r \neq 0$.
- (2) If $u < 0$, then $0 < \min \{1, -\frac{u}{v}, -\frac{u}{v} - \frac{k}{v} + 1\}$. We have $T_0 = -T$.
- (3) If $u = 0$ and $\phi \neq 1$, then $0 < \min \{1, -\frac{k}{v} + 1\}$. We have $T_0 = (\phi - 1)T$.
- (4) If $u = 0$, $\phi = 1$ and $0 < k < v$, then $-\frac{k}{v} + 1 < 1$. We have $T_0 = \frac{ck}{v} x^{k/v-1} T$.
- (5) If $u = 0$, $\phi = 1$ and $k = 0$, then $T_0 = -sx^{-1} T$. Here $s \neq 0$ by the assumption $\text{val}_\infty(g) \neq 0$.

The above calculation reveals that the valuation of the term $\Delta \cdot T$ in $(\Delta g) \cdot y_i$ is less than or equal to $\nu_\infty(T) + 1$ by Definition 6, which implies that $\text{val}_\infty(\Delta g) \leq \text{val}_\infty(g) + 1$. ■

3 Suitable bases

Let $W = (\omega_1, \dots, \omega_r)$ be a $C(x)$ -vector space basis of $A = C(x)[S]/\langle L \rangle$. Throughout this subsection, let $e_w \in C[x]$ and $M_w = (m_{i,j})_{i,j=1}^r \in C[x]^{r \times r}$ be such that

$$SW = \frac{1}{e_w} M_w W,$$

and $\gcd(e, m_{1,1}, m_{1,2}, \dots, m_{r,r}) = 1$. For algebraic function fields, the name ‘‘suitable basis’’ was introduced by Bronstein [11] in the context of lazy Hermite reduction. It refers to a basis W whose elements are (globally) integral and the denominator e_w with respect to derivation is squarefree. In this section, we seek a basis W such that W is a local integral basis at almost all orbits $\alpha + \mathbb{Z} \in \bar{C}/\mathbb{Z}$ and the denominator e_w (with respect to shift operation) is shift-free. Recall that a polynomial $p \in C[x]$ is said to be *shift-free* if $\gcd(p, \sigma^i(p)) = 1$ for all nonzero $i \in \mathbb{Z}$. So for a shift-free polynomial $p \in C[x]$, there is no $\alpha \in \bar{C}$ such that $\text{disp}_\alpha(p) > 0$.

In the differential case, every (global) integral basis is suitable. So Bronstein transforms an arbitrary basis to a suitable basis, along the way of finding an integral basis. In the shift case, since there are infinitely many singularities, even ‘‘globally’’ integral elements may not exist. Instead of finding a local integral basis at all $\alpha \in \bar{C}$, we seek for a finite subset $Z \in \bar{C}$ such that a local integral basis at Z is ‘‘suitable’’. The formal definition of suitable bases will be stated later.

We now show that the polynomial e does not depend on the choice of the basis of A but only on the $C[x]$ -submodule it generates. Let W and U be two $C(x)$ -vector space bases of A . Let $e_w, e_u \in C[x]$ and $M_w, M_u \in C[x]^{r \times r}$ be such that $SW = \frac{1}{e_w} M_w W$ and $SU = \frac{1}{e_u} M_u U$. Suppose that W and U generate the same submodule of A over $C[x]$. Then there exists a matrix $T \in C[x]^{r \times r}$ such that $W = TU$ and T is an invertible matrix over $C[x]$. Shifting both sides of the equation, we get

$$SW = \sigma(T)SU = \left(\sigma(T) \frac{1}{e_u} M_u T^{-1} \right) W = \frac{1}{e_w} M_w W.$$

Since $\sigma(T), T^{-1} \in C[x]^{r \times r}$, we have e_w divides e_u . Similarly the fact that $U = RW$ with $R = T^{-1} \in C[x]^{r \times r}$ implies that e_u divides e_w . Thus $e_w = e_u$ when e_w, e_u are monic.

For a basis $U = \{1, S, \dots, S^{r-1}\}$ of A , the denominator e_u may not be shift-free. However, we can transform U to a local integral basis W at a finite subset Z of \bar{C} such that e_w is shift-free.

Example 11. Let $T = \frac{x^2 x^1}{x+1}$, which is annihilated by $L = x^2(x+2)S - (x+1)^4 \in \mathbb{C}(x)[S]$. Then for $U = \{1\}$, we have $SU = \frac{(x+1)^4}{x^2(x+2)}U$. So $e_u = x^2(x+2)$ is not shift-free, because it has two roots -2 and 0 in the orbit \mathbb{Z} . Starting from U , we can compute a basis $W = \{(x+1)x^{-3}\}$, which is a local

integral basis at $Z = \{-1, 0\}$. The algorithm of computing an integral basis at Z guarantees that W and U generate the same $C(x)_\alpha$ -module for all $\alpha \in \mathbb{C} \setminus Z$. Then we see that $SW = xW$ and hence $e_w = 1$ is shift-free. In fact, if we choose an integer β with $\beta \geq 0$ and consider $Z = \{-1, 0, \dots, \beta\}$, then $W = \{(x+1)x^{-3} \prod_{i=1}^{\beta} (x-i)^{-1}\}$ is a local integral basis at Z and $SW = (x-\beta)W$ with $e_w = 1$ being shift-free.

In general, the above idea of finding a basis W with e_w being shift-free works as follows. Let $L = \ell_0 + \ell_1 S + \dots + \ell_r S^r \in C[x][S]$ with $\ell_0 \ell_r \neq 0$. Let $\alpha_1 + \mathbb{Z}, \dots, \alpha_I + \mathbb{Z}$ be distinct orbits such that for each i with $1 \leq i \leq I$, the product $\ell_0 \ell_r$ has at least one root in $\alpha_i + \mathbb{Z}$. Recall that for $\beta, \gamma \in \alpha + \mathbb{Z}$ with $\alpha \in \bar{C}$, we say $\beta < \gamma$ if $\beta - \gamma < 0$. For each orbit $\alpha_i + \mathbb{Z}$, let $\alpha_{i,1} < \alpha_{i,2} < \dots < \alpha_{i,J_i}$ be all roots of $\ell_0 \ell_r$ in $\alpha_i + \mathbb{Z}$. We choose an arbitrary $\beta_i \in \alpha_i + \mathbb{Z}$ such that $\alpha_{i,J_i} \leq \beta_i$. A basis W of $A = C(x)[S]/\langle L \rangle$ is called *suitable* at $\{\beta_1, \dots, \beta_I\}$ if W is a local integral basis of A at $Z = \bigcup_{i=1}^I \{\gamma \in \alpha_i + \mathbb{Z} \mid \alpha_{i,1} < \gamma \leq \beta_i\}$ and it generates the same $C(x)_\alpha$ -module as $U = \{1, S, \dots, S^{r-1}\}$ for all $\alpha \in \bar{C} \setminus Z$. So every suitable basis is a local integral basis at a finite subset Z and the set $Z \cap (\alpha_i + \mathbb{Z})$ has an upper bound β_i in $\alpha_i + \mathbb{Z}$ for each i with $1 \leq i \leq I$.

Throughout the paper, we assume that if α_{i,J_i} and $\alpha_{i',J_{i'}}$ are conjugate over C for some $1 \leq i \neq i' \leq I$, then $\beta_i - \alpha_{i,J_i} = \beta_{i'} - \alpha_{i',J_{i'}}$, which implies that β_i and $\beta_{i'}$ are also conjugate over C . Under this assumption, we show that all conjugates of β_i over C belong to $\{\beta_1, \dots, \beta_I\}$. Then a suitable basis of at $\{\beta_1, \dots, \beta_I\}$ exists and can be computed (without algebraic extension of the base field C) using the algorithm in [13] for computing an integral basis at a finite subset $Z \subseteq \bar{C}$. Since every irreducible polynomial in $C[x]$ is shift-free, all conjugates of α_{i,J_i} over C belong to distinct orbits in $\{\alpha_1 + \mathbb{Z}, \dots, \alpha_I + \mathbb{Z}\} \subseteq \bar{C}/\mathbb{Z}$. Note that if $p(x) \in C[x]$ is the minimal polynomial of $\alpha \in \bar{C}$, then $\sigma^{-k}(p(x))$ is the minimal polynomial of $\alpha + k$ for any integer k . So the number of conjugates of β_i is the same as that of α_{i,J_i} . Thus all conjugates of β_i over C belong to $\{\beta_1, \dots, \beta_I\}$.

The following theorem says that a suitable basis has the property that e_w is shift-free.

Theorem 12. *We use the above notations. Let W be a suitable basis of A at $\{\beta_1, \dots, \beta_I\}$.*

(i) *Let $e_w \in C[x]$ and $M_w \in C[x]^{r \times r}$ be such that $SW = \frac{1}{e_w} M_w W$. Then e_w is shift-free and β_1, \dots, β_I are all possible roots of e_w in \bar{C} .*

(ii) *Let $f_w \in C[x]$ and $N_w \in C[x]^{r \times r}$ be such that $W = \frac{1}{f_w} N_w SW$. Then f_w is shift-free and β_1, \dots, β_I are all possible roots of f_w in \bar{C} .*

In particular, M_w and N_w are invertible matrices over $C(x)$.

Proof. (i) A direct calculation yields that

$$SU = \begin{pmatrix} 0 & 1 & & \\ \vdots & & \ddots & \\ 0 & & & 1 \\ -\frac{\ell_0}{\ell_r} & -\frac{\ell_1}{\ell_r} & \dots & -\frac{\ell_{r-1}}{\ell_r} \end{pmatrix} U =: \frac{1}{e_u} M_u U,$$

where $e_u \in C[x]$ and $M_u \in C[x]^{r \times r}$. So every root of e_u must be a root of the leading coefficient ℓ_r . Let $T \in C(x)^{r \times r}$ be such that $W = TU$. Taking the shift operation, we get

$$SW = \sigma(T)SU = \sigma(T) \frac{1}{e_u} M_u T^{-1} W = \frac{1}{e_w} M_w W. \quad (8)$$

Since W and U generate the same $C(x)_\alpha$ -module for all $\alpha \in \bar{C} \setminus Z$, both T and T^{-1} belong to $C(x)_\alpha^{r \times r}$. So T and T^{-1} have no poles at $\bar{C} \setminus Z$, i.e., all entries of T and T^{-1} have no poles at $\alpha \in \bar{C} \setminus Z$. Then $\sigma(T)$ has no poles at $\bar{C} \setminus Z'$, where $Z' = \{\gamma - 1 \mid \gamma \in Z\} = \bigcup_{i=1}^I \{\gamma \in \alpha_i + \mathbb{Z} \mid \alpha_{i,1} \leq \gamma \leq \beta_i - 1\}$. By (8) we have

$$\begin{aligned} \{\text{roots of } e_w\} &\subseteq \{\text{poles of } \sigma(T)\} \cup \{\text{roots of } e_u\} \cup \{\text{poles of } T^{-1}\} \\ &\subseteq Z' \cup \{\text{roots of } \ell_r\} \cup Z \\ &\subseteq \bigcup_{i=1}^I (\{\alpha_{i,1}, \dots, \beta_i - 1\} \cup \{\alpha_{i,1}, \alpha_{i,2}, \dots, \alpha_{i,J_i}\} \cup \{\alpha_{i,1} + 1, \dots, \beta_i\}) \\ &\subseteq \bigcup_{i=1}^I \{\alpha_{i,1}, \alpha_{i,1} + 1, \dots, \beta_i\}; \end{aligned} \quad (9)$$

here we use the assumption that $\alpha_{i,J_i} \leq \beta_i$.

On the other hand, by Lemma 2, we know that for each $1 \leq i \leq I$, U is a local integral bases at every $\alpha \leq \alpha_{i,1}$ with $\alpha \in \alpha_i + \mathbb{Z}$. So by construction, W is a local integral basis at every $\alpha \leq \beta_i$ with $\alpha \in \alpha_i + \mathbb{Z}$. By Lemma 3, SW is a local integral basis at every $\alpha \leq \beta_i - 1$. Thus e_w has no roots in Z' because SW and W generate the same $C(x)_\alpha$ -module for all $\alpha \in Z'$. Combining this fact and the relation in (9), we get

$$\{\text{roots of } e_w\} \subseteq \{\beta_i \mid 1 \leq i \leq I\}.$$

Since $\beta_i \in \alpha_i + \mathbb{Z}$ and $\alpha_i + \mathbb{Z} \neq \alpha_j + \mathbb{Z}$ for all $1 \leq i \neq j \leq I$, it follows that e_w is shift-free.

Since both W and SW are $C(x)$ -vector space bases of A , the matrix M_w is invertible over $C(x)$.

(ii) Note that

$$U = \begin{pmatrix} -\frac{\ell_1}{\ell_0} & -\frac{\ell_2}{\ell_0} & \cdots & -\frac{\ell_r}{\ell_0} \\ 1 & & & 0 \\ & \ddots & & \vdots \\ & & 1 & 0 \end{pmatrix} SU =: \frac{1}{f_u} N_u SU,$$

where $f_u \in C[x]$ and $N_u \in C[x]^{r \times r}$. So every root of f_u must be a root of the trailing coefficient ℓ_0 . By (8), we have

$$W = T \left(\frac{1}{e_u} M_u \right)^{-1} \sigma(T)^{-1} SW = T \frac{1}{f_u} N_u \sigma(T)^{-1} SW = \frac{1}{f_w} N_w SW.$$

Using the same argument as in (i), we get

$$\begin{aligned} \{\text{roots of } f_w\} &\subseteq \{\text{poles of } T\} \cup \{\text{roots of } f_u\} \cup \{\text{poles of } \sigma(T)^{-1}\} \\ &\subseteq Z \cup \{\text{roots of } \ell_0\} \cup Z' \\ &\subseteq \bigcup_{i=1}^I \{\alpha_{i,1}, \alpha_{i,1} + 1, \dots, \beta_i\}. \end{aligned} \tag{10}$$

Since both W and SW are local integral bases at every $\alpha \leq \beta_i - 1$ with $\alpha \in \alpha_i + \mathbb{Z}$, the only possible roots of f_w are β_1, \dots, β_I . So f_w is shift-free. \blacksquare

Example 13. Let $L = (x+2)(x+3)S^2 - 2(x+2)S + 1 \in \mathbb{C}[x][S]$. It has two solutions $b_1 = \frac{1}{x!}$ and $b_2 = \frac{1}{(x+1)!}$. All roots of $\ell_0 \ell_2$ are $-3, -2 \in \mathbb{Z}$. Let $\beta = -2$ and $Z = \{-2\}$. A suitable basis of $A = \mathbb{C}(x)[S]/\langle L \rangle$ at β is $W = (1, (x+2)S)$. It is a local integral basis of A at Z and generates the same $\mathbb{C}(x)_\alpha$ -module as $U = \{1, S\}$ for all $\alpha \in \mathbb{C} \setminus Z$. Then by Theorem 12, e_w is shift-free and β is the only possible root of e_w . Indeed, we have $e_w = x+2$ and $M_w = \begin{pmatrix} 0 & 1 \\ -1 & 2 \end{pmatrix}$. Moreover, $f_w = 1$ is shift-free and $N_w = \begin{pmatrix} 2x+4 & -x-2 \\ x+2 & 0 \end{pmatrix}$.

4 Generalized Abramov-Petkovšek reduction

In this section, we use a suitable basis to decompose an element $f \in A$ into a summable part and a remainder such that the denominator of this remainder is shift-free. The lemma below presents a matrix version of Abramov-Petkovšek reduction formula in [5, Lemma 9].

Lemma 14. Let W be a suitable basis of A at $\{\beta_1, \dots, \beta_I\}$. Let $e, \tilde{e} \in C[x]$ and $M, \tilde{M} \in C[x]^{r \times r}$ be such that $SW = \frac{1}{e} MW$ and $\frac{1}{\tilde{e}} \tilde{M} = \left(\frac{1}{e} M\right)^{-1}$. Let $p_i \in C[x]$ be the minimal polynomial of β_i over C . Let $q \in C[x]$, $a \in C[x]^r$ and $\ell \in \mathbb{Z}$.

(i) If $\gcd(q, \sigma^j(p_i)) = 1$ for all $i \in \{1, \dots, I\}$ and $j \in \mathbb{Z}$, there exist $g \in A$ and $c \in C[x]^r$ such that

$$\frac{aW}{\sigma^\ell(q)} = \Delta g + \frac{cW}{qe\tilde{e}}. \tag{11}$$

(ii) If $q = p_i^m$ with $i \in \{1, \dots, I\}$ and $m > 0$, there exist $g \in A$ and $c \in C[x]^r$ such that (11) also holds.

Proof. If $\ell = 0$, we only need to take $g = 0$ and $c = e\tilde{e}a$.

If $\ell \geq 1$, we have

$$\begin{aligned} \frac{aW}{\sigma^\ell(q)} &= \frac{a}{\sigma^\ell(q)} \frac{\tilde{M}}{\tilde{e}} \frac{M}{e} W - \frac{\sigma^{-1}(a)}{\sigma^{\ell-1}(q)} \frac{\sigma^{-1}(\tilde{M})}{\sigma^{-1}(\tilde{e})} W + \frac{\sigma^{-1}(a)}{\sigma^{\ell-1}(q)} \frac{\sigma^{-1}(\tilde{M})}{\sigma^{-1}(\tilde{e})} W \\ &= \Delta \left(\frac{\sigma^{-1}(a)}{\sigma^{\ell-1}(q)} \frac{\sigma^{-1}(\tilde{M})}{\sigma^{-1}(\tilde{e})} W \right) + \frac{\sigma^{-1}(a)}{\sigma^{\ell-1}(q)} \frac{\sigma^{-1}(\tilde{M})}{\sigma^{-1}(\tilde{e})} W. \end{aligned} \quad (12)$$

By the assumption and Theorem 12.(ii), every irreducible factor of \tilde{e} is a factor of p_i for some $i \in \{1, \dots, I\}$. So in either (i) or (ii), we have $\gcd(\sigma^{\ell-1}(q), \sigma^{-1}(\tilde{e})) = 1$. By the partial fraction decomposition of rational functions, there exist $v_1, v_2 \in C[x]^r$ such that

$$\frac{\sigma^{-1}(a)}{\sigma^{\ell-1}(q)} \frac{\sigma^{-1}(\tilde{M})}{\sigma^{-1}(\tilde{e})} W = \frac{v_1 W}{\sigma^{\ell-1}(q)} + \frac{v_2 W}{\sigma^{-1}(\tilde{e})}. \quad (13)$$

Since

$$\frac{v_2 W}{\sigma^{-1}(\tilde{e})} = \Delta \left(-\frac{v_2 W}{\sigma^{-1}(\tilde{e})} \right) + \frac{\sigma(v_2)MW}{\tilde{e}e},$$

combining with (12) and (13), we have

$$\frac{aW}{\sigma^\ell(q)} \equiv \frac{v_1 W}{\sigma^{\ell-1}(q)} + \frac{\sigma(v_2)MW}{\tilde{e}e} \pmod{\Delta(A)}.$$

By the induction hypothesis, the first summand is congruent to $\frac{c_1 W}{q e \tilde{e}}$ for some $c_1 \in C[x]^r$. Setting $c = c_1 + q\sigma(v_2)M$ establishes Equation (11) in the case $\ell \geq 1$.

If $\ell \leq -1$, we have

$$\frac{aW}{\sigma^\ell(q)} = \Delta \left(-\frac{aW}{\sigma^\ell(q)} \right) + \frac{\sigma(a)MW}{\sigma^{\ell+1}(q)e}.$$

When $\ell = -1$, we take $g = -\frac{aW}{\sigma^{-1}(q)}$ and $c = \tilde{e}\sigma(a)M$. When $\ell < -1$, note that $\gcd(\sigma^{\ell+1}(q), e) = 1$ by the assumption and Theorem 12.(i). The induction can be completed in a similar way as in the case $\ell \geq 1$. \blacksquare

Recall [4, 5] that two irreducible polynomials $p, q \in C[x]$ are said to be *shift-equivalent* if $p \mid \sigma^k(q)$ for some $k \in \mathbb{Z}$, denoted by $p \sim_x q$. The relation \sim_x is an equivalence relation. By grouping together the irreducible factors in the same shift-equivalence classes, every polynomial $u \in C[x]$ has a factorization of the form

$$u = \prod_{i=1}^m \prod_{\ell=\ell_1}^{\ell_2} \sigma^\ell(q_i)^{n_{i,\ell}}$$

where $m, n_{i,\ell} \in \mathbb{N}$, $\ell_1, \ell_2 \in \mathbb{Z}$ with $\ell_1 \leq \ell_2$, $q_i \in C[x]$ is irreducible, and $q_i, q_{i'}$ are pairwise shift-inequivalent for all $1 \leq i \neq i' \leq m$. From such a factorization, applying the partial fraction decomposition of rational functions to all coefficients of $f = \frac{aW}{u} \in A$ with $a \in C[x]^r$, we can decompose f into the form

$$f = \left(f_0 + \sum_{i=1}^m \sum_{\ell=\ell_1}^{\ell_2} \frac{a_{i,\ell}}{\sigma^\ell(q_i)^{n_{i,\ell}}} \right) W \quad (14)$$

where $f_0, a_{i,j,\ell} \in C[x]^r$ and $\deg_x(a_{i,\ell}) < \deg_x(q_i^{n_{i,\ell}})$. The *degree* of a vector in $C[x]^r$ is the maximal degree of all its entries.

Proposition 15. *Let W be a suitable basis of A at $\{\beta_1, \dots, \beta_I\}$. Then any element $f \in A$ can be decomposed as*

$$f = \Delta g + h \quad \text{and} \quad h = \frac{cW}{\tilde{u}}, \quad (15)$$

where $g \in A$, $c \in C[x]^r$, $\tilde{u} \in C[x]$ and the product $\tilde{u} \prod_{i=1}^I (x - \beta_i)$ is shift-free.

Proof. We write f as in (14). Let $p_i \in C[x]$ be the minimal polynomial of β_i . If q_i is shift-equivalent to $p_{i'}$ for some $i' \in \{1, \dots, I\}$, we may choose $q_i = p_{i'}$ as a representative in this shift equivalence class. By Lemma 14, there exist $g \in A$ and $c_{i,\ell} \in C[x]^r$ such that

$$f = \Delta g + \left(f_0 + \sum_{i=1}^m \sum_{\ell=\ell_1}^{\ell_2} \frac{c_{i,\ell}}{q_i^{n_{i,\ell}} e \tilde{e}} \right) W,$$

where $SW = \frac{1}{e} MW$ and $\frac{1}{e} \tilde{M} = (\frac{1}{e} M)^{-1}$. By Theorem 12, $e \tilde{e} \prod_{i=1}^I (x - \beta_i)$ is shift-free. Then we get (15) by setting $\tilde{u} = e \tilde{e} \prod_{i=1}^m q_i^{n_i}$ with $n_i = \max_{\ell=\ell_1}^{\ell_2} \{n_{i,\ell}\}$ and $c = f_0 v + \sum_{i=1}^m \sum_{\ell=\ell_1}^{\ell_2} (v c_{i,\ell}) / (q_i^{n_{i,\ell}} e \tilde{e})$. By the assumption on suitable bases, the orbits $\beta_i + \mathbb{Z}$ with $1 \leq i \leq I$ are distinct and all conjugates of β_i belong to $\{\beta_1, \dots, \beta_I\}$. So the product $p_{i'} \prod_{i=1}^I (x - \beta_i)$ is shift-free for all $1 \leq i' \leq I$. Thus the polynomial \tilde{u} satisfies the required condition. ■

Theorem 16. *Let $h \in A$ be in (15) with $c = (c_1, \dots, c_r)$ and $\gcd(\tilde{u}, c_1, \dots, c_r) = 1$. Let $e \in C[x]$ and $M \in C[x]^{r \times r}$ be such that $SW = \frac{1}{e} MW$. If h is summable in A , then \tilde{u} divides e and $h = \Delta(bW)$ with $b \in C[x]^r$.*

Proof. Suppose h is summable in A . There exist $b = (b_1, \dots, b_r) \in C[x]^r$ and $v \in C[x]$ such that $h = \Delta(\frac{1}{v} bW)$ and $\gcd(v, b_1, \dots, b_r) = 1$. We show that v is a constant. Otherwise there exists $\alpha \in \bar{C}$ such that $\text{disp}_\alpha(v) \geq 0$. Since W is a suitable basis at $\{\beta_1, \dots, \beta_I\}$, by Proposition 5 and Theorem 12, we have that either $\text{disp}_\alpha(\tilde{u}) \geq \text{disp}_\alpha(v) + 1 \geq 1$ or $\text{disp}_\alpha((x - \beta_i)\tilde{u}) \geq 1$ for some $i \in \{1, \dots, I\}$. This contradicts the fact that $\tilde{u} \prod_{i=1}^I (x - \beta_i)$ is shift-free. Thus v is a constant in C . Then

$$h = \sigma(v^{-1}b)SW - v^{-1}bW = \left(\frac{1}{e} \sigma(v^{-1}b)M - v^{-1}b \right) W = \frac{cW}{\tilde{u}}.$$

So \tilde{u} divides e . ■

Corollary 17. *Let W be a suitable basis of A at $\{\beta_1, \dots, \beta_I\}$. Let $e \in C[x]$ and $M \in C[x]^{r \times r}$ be such that $SW = \frac{1}{e} MW$. Then any element $f \in A$ can be decomposed as*

$$f = \Delta g + \frac{1}{d} PW + \frac{1}{e} RW, \quad (16)$$

where $g \in A$, $d \in C[x]$, $P, R \in C[x]^r$ with $\deg_x(P) < \deg_x(d)$ and $d \prod_{i=1}^I (x - \beta_i)$ being shift-free. Moreover, if f is summable in A , then $P = 0$.

Proof. Let $g \in A$ and $h = \frac{cW}{\tilde{u}}$ be as in (15). Let $\{p_1, \dots, p_m\}$ be the set of minimal polynomials of the β_i 's over C for all $1 \leq i \leq I$ and let $\tilde{u} = \tilde{u} \prod_{i=1}^m p_i^{n_i}$ be such that $\tilde{u} \in C[x]$ and $\gcd(\tilde{u}, \prod_{i=1}^m p_i) = 1$. Applying the partial fraction decomposition of rational functions to all entries of c , we decompose h as

$$h = \left(h_0 + \frac{\tilde{c}}{\tilde{u}} + \sum_{i=1}^m \frac{c_i}{p_i^{n_i}} \right) W, \quad (17)$$

where $h_0, \tilde{c}, c_i \in C[x]^r$ satisfy $\deg_x(\tilde{c}) < \deg_x(\tilde{u})$ and $\deg_x(c_i) < \deg_x(p_i^{n_i})$. Let k_i be the multiplicity of p_i in e . If $n_i > k_i$, by division with remainder, we write $c_i = a_i p_i^{n_i - k_i} + b_i$ such that $a_i, b_i \in C[x]^r$ and $\deg_x(b_i) < \deg_x(p_i^{n_i - k_i})$. Then we have

$$\frac{c_i}{p_i^{n_i}} W = \left(\frac{b_i}{p_i^{n_i}} + \frac{a_i}{p_i^{k_i}} \right) W. \quad (18)$$

If $n_i \leq k_i$, then (18) holds for $b_i = 0$ and $a_i = c_i p_i^{k_i - n_i}$. Let d be the product of \tilde{u} and $p_i^{n_i}$ for all $1 \leq i \leq m$ with $b_i \neq 0$. Combining (17) and (18), we decompose h into two parts:

$$h = \left(\frac{\tilde{c}}{\tilde{u}} + \sum_{i=1}^m \frac{b_i}{p_i^{n_i}} \right) W + \left(h_0 + \sum_{i=1}^m \frac{a_i}{p_i^{k_i}} \right) W \triangleq \frac{1}{d} PW + \frac{1}{e} RW,$$

where $P = (d/\tilde{u})\tilde{c} + \sum_{i=1}^m (d/p_i^{n_i})b_i$ and $R = eh_0 + \sum_{i=1}^m (e/p_i^{k_i})a_i \in C[x]^r$. Then the product $d \prod_{i=1}^I (x - \beta_i)$ is shift-free and $\deg_x(P) < \deg_x(d)$. Since $f = \Delta g + h$, f is summable in A if and only if h is summable in A . By Theorem 12, $\{p_1, \dots, p_m\}$ are all possible irreducible factors of e . Note that for each $i \in \{1, \dots, m\}$, the multiplicity of p_i in d is either greater than k_i or equal to 0. By Theorem 16, if h is summable in A , then d belongs to C and hence $P = 0$ because $\deg_x(P) < \deg_x(d)$. ■

Example 18. Let $L \in \mathbb{C}[x][S]$ be the same operator as in Example 13 and $W = (\omega_1, \omega_2) = (1, (x+2)S)$ be a suitable basis of $A = \mathbb{C}(x)[S]/\langle L \rangle$ at $\{-2\}$. Then $e = x+2$ and $M = \begin{pmatrix} 0 & 1 \\ -1 & 2 \end{pmatrix}$. For

$$f = \frac{1}{(x+1)(x+2)}\omega_1 + \frac{x}{(x+1)(x+2)}\omega_2,$$

applying the reduction described above, we can reduce the dispersion of the denominator of f and decompose it into the form

$$f = \Delta \underbrace{\left(\frac{(-1, 1)}{x+1} W \right)}_g + \frac{1}{(x+2)^2} \underbrace{(1, -1)}_P W + \frac{1}{(x+2)} \underbrace{(-1, 2)}_R W.$$

By Theorem 16, f is not summable in A because $d \notin C$ and $P \neq 0$.

In the next section, we shall use a local integral basis at infinity to reduce the degree of S in (16).

5 Reduction at infinity

This section can be viewed as a discrete analog of Hermite reduction at infinity for D-finite functions [12]. It is also a generalization of the polynomial reduction in the hypergeometric case [14]. Let $W = (\omega_1, \dots, \omega_r)$ be a local integral basis at infinity of A . Let $\lambda \in \mathbb{Z}$ be the minimal integer such that

$$\Delta W = x^\lambda M W \quad \text{with} \quad M = (m_{i,j})_{i,j=1}^r \in C(x)_\infty^{r \times r}.$$

For convenience, we define the *degree* of a rational function $p/q \in C(x)$ as $\deg_x(p) - \deg_x(q)$, denoted by $\deg_x(p/q)$. The *degree* of a matrix in $C(x)^{r \times r}$ is the maximal degree of all its entries. With this convention, we have $\lambda = \deg_x(x^\lambda M)$. Let $f = x^k \sum_{i=1}^r a_i \omega_i \in A$ with $k \geq 0$ and $a_1, \dots, a_r \in C(x)_\infty$. In order to reduce the degree k of the coefficient vector of f , we seek $b_1, \dots, b_r, c_1, \dots, c_r \in C(x)_\infty$ such that

$$x^k \sum_{i=1}^r a_i \omega_i = \Delta \left(x^{k+1} \sum_{i=1}^r b_i \omega_i \right) + x^{k-1} \sum_{i=1}^r c_i \omega_i. \quad (19)$$

Using the relation $\Delta(uv) = \Delta(u)v + \sigma(u)\Delta(v)$ for all $u \in C(x)$ and $v \in A$, we get

$$x^k \sum_{i=1}^r a_i \omega_i = \sum_{i=1}^r \left(\Delta(b_i) x^{k+1} \omega_i + \sigma(b_i) \Delta(x^{k+1} \omega_i) + c_i x^{k-1} \omega_i \right)$$

Multiplying by x^{-k} and rewriting the difference of $x^{k+1} \omega_i$, we obtain

$$\sum_{i=1}^r a_i \omega_i = \sum_{i=1}^r \left(\Delta(b_i) x \omega_i + \sigma(b_i) x^{-k} \Delta(x^{k+1} \omega_i) + c_i x^{-1} \omega_i \right) \quad (20)$$

$$= \sum_{i=1}^r \left(\Delta(b_i) x \omega_i + \sigma(b_i) x^{-k} \Delta(x^{k+1} \omega_i) + x^{-k} \sigma(b_i x^{k+1}) x^\lambda \sum_{j=1}^r m_{i,j} \omega_j + c_i x^{-1} \omega_i \right). \quad (21)$$

Since $b_i \in C(x)_\infty$, we have $\Delta(b_i)x \in x^{-1}C(x)_\infty$. For example, $\Delta(1 + \frac{1}{x} + \dots)x = -\frac{1}{x} + \dots$. Also $x^{-k} \Delta(x^{k+1}) = (k+1) + \binom{k+1}{2} x^{-1} + \dots \equiv (k+1) \pmod{x^{-1}}$. So if $\lambda < -1$, then Equation (21) can be reduced modulo $x^{-1}C(x)_\infty$:

$$\sum_{i=1}^r a_i \omega_i \equiv \sum_{i=1}^r (k+1) \sigma(b_i) \omega_i \pmod{x^{-1}}. \quad (22)$$

It follows that $b_i \equiv \sigma^{-1}((k+1)^{-1} a_i) = (k+1)^{-1} \sigma^{-1}(a_i) \equiv (k+1)^{-1} a_i \pmod{x^{-1}}$ is the unique solution of (22) in $C(x)_\infty/\langle x^{-1} \rangle$. If $\lambda \geq -1$, then multiplying (20) by $x^{-\lambda-1}$ and reducing this equation modulo $x^{-\lambda-2}$ yields

$$\sum_{i=1}^r x^{-\lambda-1} a_i \omega_i \equiv \sum_{i=1}^r \sigma(b_i) x^{-k-\lambda-1} \Delta(x^{k+1} \omega_i) \pmod{x^{-\lambda-2}}. \quad (23)$$

Let $\psi_i := x^{-k-\lambda-1}\Delta(x^{k+1}\omega_i)$ for $i = 1, \dots, r$. To reduce the degree k , we have to show that Equation (23) always has a solution (b_1, \dots, b_r) in $(C(x)_\infty/\langle x^{-\lambda-2} \rangle)^r$. The solution $b = (b_1, \dots, b_r)$ in $(C(x)_\infty/\langle x^{-\lambda-2} \rangle)^r$ is of the form $b_i = b_{i,0} + b_{i,1}\frac{1}{x} + \dots + b_{i,\lambda+1}\frac{1}{x^{\lambda+1}}$ with $b_{i,j} \in C$. In practice, we hope that $x^{k+1}\sum_{i=1}^r b_i\omega_i$ in (19) has only polynomial coefficients and therefore assume $k \geq \max\{0, \lambda\}$.

Proposition 19. *Let $W = \{\omega_1, \dots, \omega_r\}$ be a local integral basis at infinity of A . Let $\lambda \in \mathbb{Z}$ be the minimal integer such that $\Delta(W) = x^\lambda MW$ with $M \in C(x)_\infty^{r \times r}$. For an integer $k \geq 0$, we define $\psi_i := x^{-k-\lambda-1}\Delta(x^{k+1}\omega_i)$. If $\lambda \geq -1$, then*

$$\sum_{i=1}^r C(x)_\infty\psi_i \subseteq \mathcal{O}_\infty \subseteq x^{\lambda+1}\sum_{i=1}^r C(x)_\infty\psi_i.$$

In particular, when $\lambda = -1$, we have $\sum_{i=1}^r C(x)_\infty\psi_i = \mathcal{O}_\infty$. In this case, $\{\psi_1, \dots, \psi_r\}$ forms a local integral basis at infinity.

Proof. We prove this proposition using the same technique as in [17, Lemma 10] and [12, Proposition 6]. To show $\sum_{i=1}^r C(x)_\infty\psi_i \subseteq \mathcal{O}_\infty$, we only need to show that for every $i = 1, \dots, r$, the element ψ_i is integral at infinity. Expanding ψ_i , we get $\psi_i = x^{-k-\lambda-1}\Delta(x^{k+1}\omega_i) + x^{-k-\lambda-1}\sigma(x^{k+1})\Delta(\omega_i)$. Since $\Delta(W) = x^\lambda MW$ and $M \in C(x)_\infty^{r \times r}$, it follows that the second term in ψ_i is integral at infinity. The first term in ψ_i is also integral at infinity because $\lambda + 1 \geq 0$. So ψ_i is integral at infinity.

Next we shall prove $\mathcal{O}_\infty \subseteq x^{\lambda+1}\sum_{i=1}^r C(x)_\infty\psi_i$. Suppose to the contrary that there exists an element $f \in \mathcal{O}_\infty \setminus x^{\lambda+1}\sum_{i=1}^r C(x)_\infty\psi_i$. Furthermore, we can find such an element f of the form

$$f = x^{\lambda+2}\sum_{i=1}^r \sigma(c_i)\psi_i \quad \text{with } c_i \in C(x)_\infty \text{ and } \nu_\infty(c_i) = 0 \text{ for some } i.$$

Taking $f = 0$, we shall prove that the $\{\psi_1, \dots, \psi_r\}$ is linearly independent.

Further let $g = x^2\sum_{i=1}^r \Delta(c_i)\omega_i$, which is integral at infinity for the same reason as between (21) and (22). Then also their sum

$$\begin{aligned} f + g &= x^{-k+1}\sum_{i=1}^r (\sigma(c_i)\Delta(x^{k+1}\omega_i) + \Delta(c_i)x^{k+1}\omega_i) \\ &= x^{-k+1}\sum_{i=1}^r \Delta(c_i x^{k+1}\omega_i) = x^{-k+1}\Delta(x^{k+1}h) \end{aligned}$$

must be integral, where $h = \sum_{i=1}^r c_i\omega_i$.

Since $\{\omega_1, \dots, \omega_r\}$ is a local integral basis at infinity, by [12, Lemma 2] (it also works in the setting of P-recursive sequences), we have $0 \leq \text{val}_\infty(h) < 1$. Note that

$$\text{val}_\infty(x^{k+1}h) = -k - 1 + \text{val}_\infty(h) \leq -1 + \text{val}_\infty(h) < 0;$$

here we use the assumption that $k \geq 0$, because $k = -1$ and $\text{val}_\infty(h) = 0$ imply that $\text{val}_\infty(x^{k+1}h) = 0$. Since $\text{val}_\infty(x^{k+1}h) \neq 0$, by Proposition 10 we get

$$\text{val}_\infty(x^{-k+1}\Delta(x^{k+1}h)) \leq k - 1 - k - 1 + \text{val}_\infty(h) + 1 = \text{val}_\infty(h) - 1 < 0.$$

So $x^{-k+1}\Delta(x^{k+1}h) = f + g$ is not locally integral at infinity, which contradicts the integrality of f . Hence $\mathcal{O}_\infty \subseteq x^{\lambda+1}\sum_{i=1}^r C(x)_\infty\psi_i$. ■

Note that the shift operator σ does not change the valuation ν_∞ of a rational function. The following theorem follows with the same proof as in [12, Theorem 7].

Theorem 20. *Using the same notations as in Proposition 19, let $k \geq 0$ and $\lambda \geq -1$. Then for any $a_1, \dots, a_r \in C(x)_\infty$, the linear system*

$$\sum_{i=1}^r x^{-\lambda-1}a_i\omega_i = \sum_{i=1}^r \sigma(b_i)\psi_i$$

has a solution (b_1, \dots, b_r) in $(C(x)_\infty/\langle x^{-\lambda-2} \rangle)^r$.

Example 21. Let $L \in \mathbb{C}[x][S]$ be the same operator as in Example 13. A local integral basis at infinity of $A = \mathbb{C}(x)[S]/\langle L \rangle$ is given by $\omega_1 = (x^4 + 2x^3)S - x^3 - x^2$ and $\omega_2 = (x^3 + 2x^2)S - x^2$. Then

$$\begin{pmatrix} \Delta\omega_1 \\ \Delta\omega_2 \end{pmatrix} = x^\lambda \begin{pmatrix} \frac{-x^3 - x^2 + 2x + 1}{x^2(x+2)} & 0 \\ 0 & \frac{-x^3 - x^2 + 2x + 1}{x^2(x+2)} \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}$$

with $\lambda = 0$. We apply the reduction at infinity to

$$f = \frac{x^3 + 2}{x^2(x+2)}\omega_1 + \frac{2x^3 + 3}{x^2(x+2)}\omega_2.$$

So we start with $k = 0$, $a_1 = (x^3 + 2)/(x^2(x+2))$ and $a^2 = (2x^3 + 3)/(x^2(x+2))$. Then Equation (23) leads to the following linear system for the unknowns b_1 and b_2 :

$$(x^{-1}a_1, x^{-1}a_2) \equiv (\sigma(b_1), \sigma(b_2)) \begin{pmatrix} \frac{-x^4 - x^3 + 3x^2 + 3x + 1}{x^3(x+2)} & 0 \\ 0 & \frac{-x^4 - x^3 + 3x^2 + 3x + 1}{x^3(x+2)} \end{pmatrix} \pmod{x^{-2}}.$$

By Theorem 20, this system always has a solution. Indeed, we find a solution $b_1 = -\frac{1}{x}$ and $b_2 = -\frac{2}{x}$. Then one step of the reduction at infinity simplifies f to

$$f = \Delta(-\omega_1 - 2\omega_2) + \frac{x^2 - 2x + 1}{x^2(x+2)}\omega_1 + \frac{x^2 - 2x + 2}{x^2(x+2)}\omega_2.$$

Remark 22. Let W be a local integral basis at infinity of $A = C(x)[S]/\langle L \rangle$. Let $\lambda \in \mathbb{Z}$ be the minimal integer such that $\Delta W = x^\lambda MW$ with $M \in C(x)_\infty^{r \times r}$. If the operator L (of order r) admits r linearly independent solutions in

$$\bar{C}[[x^{-1}]] := \bigcup_{\alpha \in \bar{C}} x^\alpha \bar{C}[[x^{-1/v}]][\log(x)],$$

then $\lambda \leq -1$. This is similar to the Fuchsian D -finite case, see [17, Lemma 4].

Let $W = \{\omega_1, \dots, \omega_r\}$ be a local integral basis at infinity. Let $\lambda \in \mathbb{Z}$ be the minimal integer such that $\Delta W = x^\lambda MW$ with $M \in C(x)_\infty^{r \times r}$. By a repeated application of the reduction at infinity, one can reduce the degree in x as far as possible and decompose $f \in A$ as

$$f = \Delta g + h \quad \text{with} \quad h = \sum_{i=1}^r h_i \omega_i \tag{24}$$

where $g \in A$, $h_i \in C(x)$ with $\deg_x(h_i) < \max\{0, \lambda\}$ for all i and the coefficients of g are polynomials. The following lemma follows from Proposition 10 with the same proof as in [12, Lemma 10].

Lemma 23. Let $h \in A$ be as in (24). If h is summable in A , then $h = \Delta(\sum_{i=1}^r b_i \omega_i)$ with $b_i \in C(x)$ and $\deg_x(b_i) \leq \max\{0, \lambda\}$ for all $i \in \{1, \dots, r\}$.

6 An additive decomposition

Now we combine the generalized Abramov-Petkovšek reduction and the reduction at infinity to decompose a P -recursive sequence f as $f = \Delta g + h$ such that f is summable if and only if the remainder h is zero. This procedure is similar to the hypergeometric case [14], P -recursive case [10, 31] and the differential case [8, 9, 12, 15, 17, 32]. It provides an alternative method for solving the accurate integration problem for P -recursive sequences [3].

In this section, we need two bases to perform the generalized Abramov-Petkovšek reduction and the reduction at infinity, respectively. Let $W = (\omega_1, \dots, \omega_r)$ be a suitable basis of A at $\{\beta_1, \dots, \beta_I\}$ that is normal at infinity. By the same proof as in [17, Lemma 18], there exists $T = \text{diag}(x^{\tau_1}, \dots, x^{\tau_r}) \in C(x)^{r \times r}$ with $\tau_i \in \mathbb{Z}$ such that $V := TW$ is a local integral basis at infinity. Let $e, a \in C[x]$ and $M, B \in C[x]^{r \times r}$ be such that $SW = \frac{1}{e}MW$ and $\Delta V = \frac{1}{a}BV$. Since the difference of V is $\Delta V = (S - 1)TW = (\frac{1}{e}\sigma(T)M - T)T^{-1}V$, we may assume that $a = x^{\lambda_1}(x+1)^{\lambda_2}e$ for some $\lambda_1, \lambda_2 \in \mathbb{N}$. For $\mu, \delta \in \mathbb{Z}$ with $\mu \leq \delta$, we define a subspace of Laurent polynomials in $C[x, x^{-1}]$ as $C[x]_{\mu, \delta} := \{\sum_{i=\mu}^{\delta} a_i x^i \mid a_i \in C\}$. The following theorem decomposes any $f \in A$ into a summable part and a remainder such that the remainder belongs to a finite dimensional vector space over C .

Theorem 24. Let $W, V \in A^r$ be as described above. Then any element $f \in A$ can be decomposed into

$$f = \Delta g + \frac{1}{d}PW + \frac{1}{a}QV, \quad (25)$$

where $g \in A$, $d \in C[x]$, $P \in C[x]^r$, $Q \in C[x]_{\mu, \delta}^r$ with $\deg_x(P) < \deg_x(d)$, $\mu = \min\{-\tau_1, \dots, -\tau_r, 0\}$, $\delta = \max\{\deg_x(a), \deg_x(B)\} - 1$ and the product $d \prod_{i=1}^I (x - \beta_i)$ being shift-free. Moreover, f is summable in A if and only if $R = 0$ and

$$\frac{1}{a}QV \in \Delta(U) \quad \text{with} \quad U = \{cV \mid c \in C[x]_{\mu, \delta'}^r\},$$

where $\delta' = \max\{0, \deg_x(B) - \deg_x(a)\}$.

Proof. After performing the generalized Abramov-Petkovšek reduction in Section 4, it follows from Corollary 17 that

$$f = \Delta g + \frac{1}{d}PW + \frac{1}{e}RW, \quad (26)$$

where $g \in A$, $d \in C[x]$, $P, R \in C[x]^r$ with $\deg_x(P) < \deg_x(d)$ and $d \prod_{i=1}^I (x - \beta_i)$ being shift-free. We rewrite the last summand in (26) with respect to the basis V :

$$\frac{1}{e}RW = \frac{1}{a}\tilde{R}V,$$

where $\tilde{R} = x^{\lambda_1}(x+1)^{\lambda_2}RT^{-1} \in x^\mu C[x]^r$. Since V is a local integral basis at infinity, using the reduction at infinity in Section 5, we obtain from (24) that

$$\frac{1}{e}RW = \Delta(R_1V) + \frac{1}{a}R_2V, \quad (27)$$

where $R_1 \in C[x]^r$ and $R_2 \in x^\mu C[x]^r$ satisfies

$$\deg_x\left(\frac{R_2}{a}\right) \leq \max\{0, \deg_x\left(\frac{B}{a}\right)\} - 1.$$

This implies that $\deg_x(R_2) \leq \max\{\deg_x(a), \deg_x(B)\} - 1 = \delta$. Thus $R_2 \in C[x]_{\mu, \delta}^r$ and we finally obtain the decomposition (25) by setting $g = \tilde{g} + R_1V$ and $Q = R_2$.

For the last assertion, assume that f is summable (the other direction of the equivalence holds trivially). Then Corollary 17 implies that $P = 0$. Hence the last summand in (26) is also summable. By Theorem 16, there exists $b \in C[x]^r$ such that $\frac{1}{e}RW = \Delta(bW)$. By Equation (27), we get

$$\frac{1}{a}QV = \frac{1}{e}RW - \Delta(R_1V) = \Delta((bT^{-1} - R_1)V) = \Delta(cV),$$

where $c = bT^{-1} - R_1 \in x^\mu C[x]^r$. Since V is a local integral basis at infinity, it follows from Lemma 23 that $\deg_x(c) \leq \max\{0, \deg_x\left(\frac{B}{a}\right)\} = \delta'$. Thus $c \in C[x]_{\mu, \delta'}^r$. ■

The remaining step is to reduce all summable P-recursive sequences to zero. Note that in Theorem 24, U is a C -vector space of dimension $r(\delta' - \mu' + 1)$. Since Δ is C -linear, it follows that $\Delta(U)$ is also a finite-dimensional C -vector space. Using Gaussian elimination, f in (25) can be further decomposed as

$$f = \Delta\tilde{g} + \frac{1}{d}PW + \frac{1}{a}Q_2V, \quad (28)$$

where $\tilde{g} = g + g_1$ with $g_1 \in U$ and $Q_2 \in C[x]_{\mu, \delta}^r$ such that f is summable in A if and only if $P = Q_2 = 0$. This decomposition (28) is called an *additive decomposition* of f .

Proposition 25. We use the same notations as in Theorem 24. The vector Q_2 in (28) belongs to a C -vector space of dimension at most

$$r(\delta - \mu + 1) = \max\{\deg_x(a), \deg_x(B)\} + \max\{\tau, 0\},$$

where $\tau = \max\{\tau_1, \dots, \tau_r\}$.

7 Creative Telescoping

Let $K(x)[S_t, S_x]$ with $K = C(t)$ be an Ore algebra, in which S_t and S_x are the shift operators with respect to t and x , respectively. Let J be a left ideal of $K(x)[S_t, S_x]$ such that $A = K(x)[S_t, S_x]/J$ is a $K(x)$ -vector space of dimension r . Assume that there exists a cyclic vector γ with respect to S_x . This means that $\{\gamma, S_x\gamma, \dots, S_x^{r-1}\gamma\}$ is a basis of A over $K(x)$. Then γ is annihilated by L and $S_t - u_t$ for some $L, u_t \in K(x)[S_x]$. We further assume that $L = \ell_0 + \ell_1 S_x + \dots + \ell_r S_x^r \in K[x][S_x]$ with $\ell_0 \ell_r \neq 0$. Every element f in A can be uniquely written as $P_f \gamma + J$ for some $P_f \in K(x)[S_x]$. The map sending f to $P_f + \langle L \rangle$ gives an isomorphism from A to $K(x)[S_x]/\langle L \rangle$ as a $K(x)[S_x]$ -module. Using this isomorphism, for any $f \in A$, we can apply our additive decomposition to test whether f is summable (in x). If $f \in A$ is not summable, one can ask to find a nonzero operator $T \in C(t)[S_t]$ (free of x) such that $T(f)$ is summable. Such an operator T if it exists is called a *telescoper* for f . Applying the additive decomposition with respect to x in Section 6 to $S_t^i f \in A$ yields that

$$S_t^i f = \Delta_x(g_i) + h_i \quad \text{with} \quad h_i = \frac{1}{d} P_i W + \frac{1}{a} Q_i V \quad (29)$$

where $g_i \in A$, $d \in K[x]$, $P_i \in K[x]^r$, $Q_i \in K[x]_{\mu, \delta}^r$ with $\deg_x(P_i) < \deg_x(d)$. Moreover, $S_t^i f$ is summable in A if and only if $h_i = 0$. To compute telescopers for bivariate P-recursive sequences, we are confronted with the same problem as in the hypergeometric case [14]. The sum of two remainders in the additive decomposition may not be a remainder. Using the same technique as in the hypergeometric case [14, Section 5], one can replace h_i by \tilde{h}_i such that the linear combination of the \tilde{h}_i 's over $C(t)$ is still a remainder. We shall explain how it works by the following example. If there exist $c_0, c_1, \dots, c_m \in C(t)$ such that $\sum_{i=0}^m c_i \tilde{h}_i = 0$, then $T = \sum_{i=0}^m c_i S_t^i$ is a telescoper for f . This approach is the method of reduction-based telescoping and was developed for various classes of functions [7–10, 14, 15, 17, 31, 32].

Example 26. Let $F = x + t^2 + \frac{1}{x+1}$, which is annihilated by

$$L = (x+2)(x^2 + (t^2+1)x+1)S_x^2 - (x^3 + (t^2+5)x^2 + (3t^2+7)x + t^2+4)S_x + x^2 + (t^2+3)x + t^2 + 3$$

and

$$S_t - \frac{(2t+1)(x+1)}{x^2 + (t^2+1)x+1} S_x - \frac{x^2 + (t^2+2)x - 2t}{x^2 + (t^2+1)x+1}.$$

A suitable basis of $A = K(x)[S_x]/\langle L \rangle$ with $K = \mathbb{C}(t)$ at $\{\beta_1, \beta_2, \beta_3\}$ is

$$W = (\omega_1, \omega_2) = \left(\frac{(t^2-1)}{x^2 + (t^2+1)x+1} ((x+t^2)S_x - (x+t^2+1)), \frac{1}{x^2 + (t^2+1)x+1} (S_x - \frac{x+t^2}{t^2-1}) \right),$$

where $\beta_1 = -2$, β_2 and β_3 are distinct roots of $x^2 + (t^2+1)x+1$. A local integral basis at infinity is $V = (v_1, v_2) = (x\omega_1, \omega_2)$. Then

$$\begin{aligned} \begin{pmatrix} S_x \omega_1 \\ S_x \omega_2 \end{pmatrix} &= \frac{1}{x+2} \begin{pmatrix} 1 & 0 \\ -\frac{x+1}{(t^2-1)^2} & x+2 \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}, & \begin{pmatrix} S_t \omega_1 \\ S_t \omega_2 \end{pmatrix} &= \begin{pmatrix} \frac{t(t+2)}{t^2-1} & 0 \\ 0 & \frac{t^2-1}{t(t+2)} \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \end{pmatrix}, \\ \begin{pmatrix} \Delta_x v_1 \\ \Delta_x v_2 \end{pmatrix} &= \frac{1}{x(x+2)} \begin{pmatrix} -x^2 - x + 1 & 0 \\ -\frac{x+1}{(t^2-1)^2} & 0 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}, & \begin{pmatrix} S_t v_1 \\ S_t v_2 \end{pmatrix} &= \begin{pmatrix} \frac{t(t+2)}{t^2-1} & 0 \\ 0 & \frac{t^2-1}{t(t+2)} \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}. \end{aligned}$$

We compute a minimal telescoper for $\frac{1}{x+t}F$, which corresponds to $f = \frac{1}{x+t} \in A$. Its representation in the bases is

$$\begin{aligned} f &= \frac{1}{x+t} (\omega_1 - (t^2-1)(x+t^2)\omega_2) \\ &= \frac{1}{x+t} (1, -(t-1)^2 t(t+1)) W + (0, -(t^2-1)) V. \end{aligned}$$

After the reduction at infinity, we obtain

$$f = \Delta_x \left(\left(\frac{1}{t^2-1}, -(t^2-1)x \right) V \right) + \frac{1}{x+t} (1, -(t-1)^2 t(t+1)) W + \frac{1}{x(x+2)} \left(-\frac{1}{t^2-1} (x+2), 0 \right) V.$$

Since $a = x(x+2)$, $\mu = -1$, $\delta = 1$ and $\delta' = 0$, a basis of the space of all summable sequences in the form $\frac{1}{a} QV$ with $Q \in K[x]_{\mu, \delta}^2$ is given by $\Delta_x \left(\frac{1}{x} v_1 \right) = -\frac{x+1}{x(x+2)} v_1$. An additive decomposition of f is

$$f = \Delta_x(g_0) + h_0 \quad \text{with} \quad h_0 = \frac{1}{x+t} (1, -(t-1)^2 t(t+1)) W + \frac{1}{x(x+2)} \left(-\frac{1}{t^2-1}, 0 \right) V,$$

where $g_0 = (\frac{1}{t^2-1}, -(t^2-1)x)V + (\frac{1}{(t^2-1)x}, 0)V$. Set $\tilde{h}_0 = h_0$. Next we consider $S_t f$, which has an additive decomposition

$$S_t f = \Delta_x(S_t g_0) + h_1 \quad \text{with} \quad h_1 = \frac{1}{x+t+1}(\frac{t(t+2)}{t^2-1}, -(t-1)t(t+1)^2)W + \frac{1}{x(x+2)}(-\frac{1}{t^2-1}, 0)V.$$

The sum of two remainders h_0 and h_1 is not a remainder, because $(x+t)(x+t+1)$ is not shift-free with respect to x . Applying the generalized Abramov-Petkovšek reduction to h_1 by the formula (12), we get

$$S_t f = \Delta_x(\dots) + \frac{1}{x+t}(\frac{t}{t^2-1}(x+t+2), -(t-1)t(t+1)^2)W + \frac{1}{x(x+2)}(-\frac{1}{t^2-1}, 0)V,$$

which has another additive decomposition

$$S_t f = \Delta_x(\tilde{g}_1) + \tilde{h}_1 \quad \text{with} \quad \tilde{h}_1 = \frac{1}{x+t}(\frac{2t}{t^2-1}, -(t-1)t(t+1)^2)W + \frac{1}{x(x+2)}(\frac{1}{t+1}, 0)V,$$

where $\tilde{g}_1 \in A$. Now the sum of \tilde{h}_0 and \tilde{h}_1 is a remainder. Since \tilde{h}_0 and \tilde{h}_1 are linearly independent over $\mathbb{C}(t)$, we continue with $S_t^2 f$, $S_t^3 f$ and compute their refined additive decompositions $S_t^i(f) = \Delta_x(\tilde{g}_i) + \tilde{h}_i$ for $i = 2, 3$, where $\tilde{g}_2, \tilde{g}_3 \in A$ and

$$\begin{aligned} \tilde{h}_2 &= \frac{1}{x+t}(\frac{t^2+2}{t-1}, -(t-1)(t+1)^2(t+2))W + \frac{1}{x(x+2)}(-\frac{t^2}{t^2-1}, 0)V, \\ \tilde{h}_3 &= \frac{1}{x+t}(-\frac{(t+2)(t^3-2t-3)}{t^2-1}, -(t+2)(t+3)(t^2-1))W + \frac{1}{x(x+2)}(\frac{t^2+t-1}{t-1}, 0)V. \end{aligned}$$

Now we see the remainders $\tilde{h}_0, \tilde{h}_1, \tilde{h}_2, \tilde{h}_3$ are linearly dependent over $\mathbb{C}(t)$, which gives rise to a minimal telescoper $(3t^2 + 3t + 2)S_t^3 + (3t^3 + 3t^2 - 4t - 6)S_t^2 - (6t^3 + 15t^2 + 13t + 2)S_t + 3t^3 + 9t^2 + 8t$.

References

- [1] Sergei A. Abramov. On the summation of rational functions. *USSR Computational Mathematics and Mathematical Physics*, 11(4):324–330, 1971.
- [2] Sergei A. Abramov, Manuel Bronstein, Marko Petkovšek, and Carsten Schneider. On rational and hypergeometric solutions of linear ordinary difference equations in $\Pi\Sigma^*$ -field extensions. *Journal of Symbolic Computation*, 107:23–66, 2021.
- [3] Sergei A. Abramov and Mark van Hoeij. Integration of solutions of linear functional equations. *Integral Transforms and Special Functions*, 8(1-2):3–12, 1999.
- [4] Sergei A. Abramov and Marko Petkovšek. Minimal decomposition of indefinite hypergeometric sums. In *Proceedings of the 2001 International Symposium on Symbolic and Algebraic Computation*, page 7–14, New York, NY, USA, 2001. ACM.
- [5] Sergei A. Abramov and Marko Petkovšek. Rational normal forms and minimal decompositions of hypergeometric terms. *Journal of Symbolic Computation*, 33(5):521–543, 2002.
- [6] Shayea Aldossari. *Algorithms for Simplifying Differential Equations*. PhD thesis, Florida State University, 2020.
- [7] Alin Bostan, Shaoshi Chen, Frédéric Chyzak, and Ziming Li. Complexity of creative telescoping for bivariate rational functions. In *Proceedings of the 2010 International Symposium on Symbolic and Algebraic Computation*, page 203–210, New York, NY, USA, 2010. ACM.
- [8] Alin Bostan, Shaoshi Chen, Frédéric Chyzak, Ziming Li, and Guoce Xin. Hermite reduction and creative telescoping for hyperexponential functions. In *Proceedings of the 2013 International Symposium on Symbolic and Algebraic Computation*, pages 77–84, New York, NY, USA, 2013. ACM.
- [9] Alin Bostan, Frédéric Chyzak, Pierre Lairez, and Bruno Salvy. Generalized Hermite reduction, creative telescoping and definite integration of D-finite functions. In *Proceedings of the 2018 International Symposium on Symbolic and Algebraic Computation*, pages 95–102, New York, NY, USA, 2018. ACM.

- [10] Hadrien Brochet and Bruno Salvy. Reduction-based creative telescoping for definite summation of D-finite functions. *Preprint on arXiv: 2307.07216*, 2023.
- [11] Manuel Bronstein. The lazy Hermite reduction. Technical Report 3562, INRIA, 1998.
- [12] Shaoshi Chen, Lixin Du, and Manuel Kauers. Hermite reduction for D-finite functions via integral bases. In *Proceedings of the 2023 International Symposium on Symbolic and Algebraic Computation*, page 155–163, New York, NY, USA, 2023. ACM.
- [13] Shaoshi Chen, Lixin Du, Manuel Kauers, and Thibaut Verron. Integral bases for P-recursive sequences. In *Proceedings of the 2020 International Symposium on Symbolic and Algebraic Computation*, page 91–98, New York, NY, USA, 2020. ACM.
- [14] Shaoshi Chen, Hui Huang, Manuel Kauers, and Ziming Li. A modified Abramov-Petkovšek reduction and creative telescoping for hypergeometric terms. In *Proceedings of the 2015 International Symposium on Symbolic and Algebraic Computation*, pages 117–124, New York, NY, USA, 2015. ACM.
- [15] Shaoshi Chen, Manuel Kauers, and Christoph Koutschan. Reduction-based creative telescoping for algebraic functions. In *Proceedings of the 2016 International Symposium on Symbolic and Algebraic Computation*, pages 175–182, New York, NY, USA, 2016. ACM.
- [16] Shaoshi Chen and Michael F. Singer. Residues and telescopers for bivariate rational functions. *Advances in Applied Mathematics*, 49(2):111–133, 2012.
- [17] Shaoshi Chen, Mark van Hoeij, Manuel Kauers, and Christoph Koutschan. Reduction-based creative telescoping for fuchsian D-finite functions. *Journal of Symbolic Computation*, 85:108–127, 2018.
- [18] Frédéric Chyzak. An extension of Zeilberger’s fast algorithm to general holonomic functions. *Discrete Mathematics*, 217:115–134, 2000.
- [19] Frédéric Chyzak, Manuel Kauers, and Bruno Salvy. A non-holonomic systems approach to special function identities. In *Proceedings of ISSAC’09*, pages 111–118, 2009.
- [20] Frédéric Chyzak and Bruno Salvy. Non-commutative elimination in Ore algebras proves multivariate identities. *Journal of Symbolic Computation*, 26:187–227, 1998.
- [21] Ralph William Gosper, Jr. Decision procedure for indefinite hypergeometric summation. *Proceedings of the National Academy of Sciences*, 75(1):40–42, 1978.
- [22] Manuel Kauers. *D-finite Functions*. Springer Cham, 2023.
- [23] Manuel Kauers and Christoph Koutschan. Integral D-finite functions. In *Proceedings of the 2015 International Symposium on Symbolic and Algebraic Computation*, pages 251–258, New York, NY, USA, 2015. ACM.
- [24] Christoph Koutschan. Creative telescoping for holonomic functions. In *Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions*, pages 171–194. Springer, 2013.
- [25] Peter Paule. Greatest factorial factorization and symbolic summation. *Journal of Symbolic Computation*, 20(3):235–268, 1995.
- [26] Marko Petkovšek, Herbert S. Wilf, and Doron Zeilberger. *A = B*. A. K. Peters Ltd., Wellesley, MA, 1996.
- [27] Clemens Raab. *Definite Integration in Difference Fields*. PhD thesis, Johannes Kepler University, 2012.
- [28] Carsten Schneider. A difference ring theory for symbolic summation. *Journal of Symbolic Computation*, 72(1–2):82–127, 2016.
- [29] Carsten Schneider. Summation theory II: Characterizations of $RII\Sigma^*$ -extensions and algorithmic aspects. *Journal of Symbolic Computation*, 80(3):616–664, 2017.
- [30] Barry M. Trager. *Integration of Algebraic Functions*. PhD thesis, Massachusetts Institute of Technology, 1984.

- [31] Joris van der Hoeven. Creative telescoping using reductions. hal-01773137, 2018.
- [32] Joris van der Hoeven. Constructing reductions for creative telescoping: the general differentially finite case. *Applicable Algebra in Engineering, Communication and Computing*, 32(5):575–602, 2021.
- [33] Mark van Hoeij. An algorithm for computing an integral basis in an algebraic function field. *Journal of Symbolic Computation*, 18(4):353–363, 1994.
- [34] Doron Zeilberger. A holonomic systems approach to special functions identities. *Journal of Computational and Applied Mathematics*, 32:321–368, 1990.