



ELSEVIER

Contents lists available at ScienceDirect

Journal of Symbolic Computation

www.elsevier.com/locate/jsc



Desingularization of Ore operators

Shaoshi Chen^{a,1}, Manuel Kauers^{b,2}, Michael F. Singer^{c,3}^a KLMM, AMSS, Chinese Academy of Sciences, 100190 Beijing, China^b Institute for Algebra, Johannes Kepler University, Linz, Austria^c Department of Mathematics, North Carolina State University, Raleigh, NC, USA

ARTICLE INFO

Article history:

Received 14 October 2014

Accepted 12 October 2015

Available online 9 November 2015

Keywords:

D-finite functions

Apparent singularities

Computer algebra

Ore operators

ABSTRACT

We show that Ore operators can be desingularized by calculating a least common left multiple with a random operator of appropriate order, thereby turning a heuristic used for many years in several computer algebra systems into an algorithm. Our result can be viewed as a generalization of a classical result about apparent singularities of linear differential equations.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Consider a linear ordinary differential equation, like for example

$$x(1-x)f'(x) - f(x) = 0.$$

The leading coefficient polynomial $x(1-x)$ of the equation is of special interest because every point ξ which is a singularity of some solution of the differential equation is also a root of this polynomial. However, the converse is in general not true. In the example above, the root $\xi = 1$ indicates the singularity of the solution $x/(1-x)$, but there is no solution which has a singularity at the other root $\xi = 0$. To see this, observe that after differentiating the equation, we can cancel (“remove”) the factor x from it. The result is the higher order equation

$$(1-x)f''(x) - 2f'(x) = 0,$$

E-mail addresses: schen@amss.ac.cn (S. Chen), manuel.kauers@jku.at (M. Kauers), singer@math.ncsu.edu (M.F. Singer).

¹ Supported by NSFC grant 11501552 and a 973 project (2011CB302401) and the President Fund of Academy of Mathematics and Systems Science, CAS (2014-cjrwlxz-chshsh).

² Supported by FWF grants Y464-N18, F5004.

³ Supported by National Science Foundation Grant CCF-1017217.

<http://dx.doi.org/10.1016/j.jsc.2015.11.001>

0747-7171/© 2015 Elsevier Ltd. All rights reserved.

whose solution space contains the solution space of the original equation. Such a calculation is called *desingularization*. The factor x is said to be *removable*.

Given a differential equation, it is of interest to decide which factors of its leading coefficient polynomial are removable, and to construct a higher order equation in which all the removable factors are removed. A classical algorithm, which is known since the end of the 19th century (Schlesinger, 1895; Ince, 1926), proceeds by taking the least common left multiple of the given differential operator with a suitably constructed auxiliary operator. This algorithm is summarized in Section 2 below. At the end of the 20th century, the corresponding problem for linear recurrence equations was studied and algorithms for identifying removable factors have been found and their relations to “singularities” of solutions have been investigated (Abramov and van Hoeij, 1999, 2003; Abramov et al., 2006). Also some steps towards a unified theory for desingularization of Ore operators have been made (Chyzak et al., in preparation; Chen et al., 2013). Possible connections to Ore closures of an operator ideal have been noted in Chyzak et al. (in preparation) and within the context of order-degree curves (Chen et al., 2013, Chen and Kauers, 2012a; 2012b). These will be further developed in a future paper.

Our contribution in the present article is a three-fold generalization of the classical desingularization algorithm for differential equations. Our main result (Theorem 6 below) says that (a) instead of the particular auxiliary operator traditionally used, almost every other operator of appropriate order also does the job, (b) also the case is covered where a multiple root can't be removed completely but only its multiplicity can be reduced, and (c) the technique works not only for differential operators but for every Ore algebra. Code fragments in the Maple library (e.g., the function DEtools/Homomorphisms/AppCheck) indicate that some people have already observed before us that taking lclm with a random operator tends to remove removable factors and used this as a heuristic. We give here for the first time a rigorous justification of this phenomenon.

For every removable factor p there is a smallest $n \in \mathbb{N}$ such that removing p from the operator requires increasing the order of the operator by at least n . Classical desingularization algorithms compute for each factor p an upper bound for this n , and then determine whether or not it is possible to remove p at the cost of increasing the order of the operator by at most n . In the present paper, we do not address the question of finding bounds on n but only discuss the second part: assuming some $n \in \mathbb{N}$ is given as part of the input, we consider the task of removing as many factors as possible without increasing the order of the operator by more than n . Of course, for Ore algebras where it is known how to obtain bounds on n , these bounds can be combined with our result.

Recall the notion of Ore algebras (Ore, 1933). Let K be a field of characteristic zero. Let $\sigma : K[x] \rightarrow K[x]$ be a ring automorphism that leaves the elements of K fixed, and let $\delta : K[x] \rightarrow K[x]$ be a K -linear map satisfying the law $\delta(uv) = \delta(u)v + \sigma(u)\delta(v)$ for all $u, v \in K[x]$. The algebra $K[x][\partial]$ consists of all polynomials in ∂ with coefficients in $K[x]$ together with the usual addition and the unique (in general noncommutative) multiplication satisfying $\partial u = \sigma(u)\partial + \delta(u)$ for all $u \in K[x]$ is called an *Ore algebra*. The field K is called the *constant field* of the algebra. Every nonzero element L of an Ore algebra $K[x][\partial]$ can be written uniquely in the form

$$L = \ell_0 + \ell_1\partial + \cdots + \ell_r\partial^r$$

with $\ell_0, \dots, \ell_r \in K[x]$ and $\ell_r \neq 0$. We call $\deg_{\partial}(L) := r$ the *order* of L and $\text{lc}_{\partial}(L) := \ell_r$ the *leading coefficient* of L . Roots of the leading coefficient ℓ_r are called *singularities* of L . Prominent examples of Ore algebras are the algebra of linear differential operators (with $\sigma = \text{id}$ and $\delta = \frac{d}{dx}$; we will write D instead of ∂ in this case) and the algebra of linear recurrence operators (with $\sigma(x) = x + 1$ and $\delta = 0$; we will write S instead of ∂ in this case).

We shall suppose that the reader is familiar with these definitions and facts, and will make free use of well-known facts about Ore algebras, as explained, for instance, in Ore (1933), Bronstein and Petkovšek (1996), Abramov et al. (2005). In particular, we will make use of the notion of least common left multiples (lclm) of elements of Ore algebras: $L \in K(x)[\partial]$ is a *common left multiple* of $P, Q \in K(x)[\partial]$ if we have $L = UP = VQ$ for some $U, V \in K(x)[\partial]$, it is called a *least common left multiple* if there is no common left multiple of lower order. Least common left multiples are unique up to left-multiplication by nonzero elements of $K(x)$. By $\text{lclm}(P, Q)$ we denote a least common left multiple whose coefficients belong to $K[x]$ and share no common divisors in $K[x]$. Note that $\text{lclm}(P, Q)$ is

unique up to (left-)multiplication by nonzero elements of K . Efficient algorithms for computing least common left multiples are available (Bostan et al., 2012).

2. The differential case

In order to motivate our result, we begin by recalling the classical results concerning the desingularization of linear differential operators. See the appendix of Abramov et al. (2006) for further details on this case.

Let $L = \ell_0 + \ell_1 D + \dots + \ell_r D^r \in K[x][D]$ be a differential operator of order r . Consider the power series solutions of L . It can be shown that $x \nmid \ell_r$ if and only if L admits r power series solutions with valuation α , for $\alpha = 0, \dots, r - 1$. Therefore, if $x \mid \ell_r$, then this factor is removable if and only if there exists some left multiple M of L , say with $\text{deg}_\partial(M) = s$, such that M admits a power series solution with valuation α for every $\alpha = 0, \dots, s - 1$. This is the case if and only if L has r linearly independent power series solutions with integer exponents $0 \leq \alpha_1 < \alpha_2 < \dots < \alpha_r$, because in this case (and only in this case) we can construct a left multiple M of L with power series solutions $x^\alpha + \dots$ for each $\alpha = 0, \dots, \max\{\alpha_1, \dots, \alpha_r\} - 1$, by adding power series of the missing valuations to the solution space of L .

These observations suggest the following desingularization algorithm for operators $L \in K[x][\partial]$ with $x \mid \text{lc}_\partial(L)$. First find the set $\{\alpha_1, \dots, \alpha_\ell\} \subseteq \mathbb{N}$ of all exponents α_i for which there exist power series solutions $x^{\alpha_i} + \dots$. If $\ell < r$, return “not desingularizable” and stop. Otherwise, let $m = \max\{\alpha_1, \dots, \alpha_\ell\}$ and let $e_1, e_2, \dots, e_{m-\ell}$ be those nonnegative integers which are at most m but not among the α_i . Return the operator

$$M = \text{lclm}(L, A),$$

where

$$A := \text{lclm}(xD - e_1, xD - e_2, \dots, xD - e_{m-\ell}).$$

Note that among the solutions of A there are the monomials $x^{e_1}, x^{e_2}, \dots, x^{e_{m-\ell}}$, and that the solutions of M are linear combinations of solutions of A and solutions of L . Therefore, by the choice of the e_j and the remarks made above, M is desingularized.

Example 1. Consider the operator

$$\begin{aligned} L &= (x - 1)(x^2 - 3x + 3)x D^2 - (x^2 - 3)(x^2 - 2x + 2)D \\ &\quad + (x - 2)(2x^2 - 3x + 3) \in K[x][D]. \end{aligned}$$

This operator has power series solutions with minimal exponents $\alpha = 0$ and $\alpha = 3$. Their first terms are

$$\begin{aligned} &1 + x + \frac{1}{2}x^2 - \frac{1}{8}x^4 - \frac{19}{120}x^5 - \frac{119}{720}x^6 + \dots, \\ &x^3 + x^4 + x^5 + x^6 + \dots. \end{aligned}$$

The missing exponents are $e_1 = 1$ and $e_2 = 2$. Therefore we take

$$A := \text{lclm}(xD - 1, xD - 2) = x^2 D^2 - 2xD + 2$$

and calculate

$$\begin{aligned} M = \text{lclm}(L, A) &= (x^5 - 2x^4 + 4x^3 - 9x^2 + 12x - 6)D^4 \\ &\quad - (x^5 - 2x^4 + x^3 - 12x^2 + 24x - 24)D^3 \\ &\quad - (3x^3 + 9x^2)D^2 + (6x^2 + 18x)D - (6x + 18). \end{aligned}$$

Note that we have $x \nmid \text{lc}_\partial(M)$, as predicted.

In the form sketched above, the algorithm applies only to the singularity 0. In order to get rid of a different singularity, move this singularity to 0 by a suitable change of variables, then proceed as described above, and after that undo the change of variables. Note that by removing the singularity 0 we will in general introduce new singularities at other points.

3. Removable factors

We now turn from the algebra of linear differential operators to arbitrary Ore algebras. In the general case, removability of a factor of the leading coefficient is defined as follows.

Definition 2. Let $L \in K[x][\partial]$ and let $p \in K[x]$ be such that $p \mid \text{lc}_\partial(L) \in K[x]$. We say that p is *removable* from L at order n if there exist some $P \in K(x)[\partial]$ with $\text{deg}_\partial(P) = n$ and some $v, w \in K[x]$ with $\text{gcd}(p, w) = 1$ such that $PL \in K[x][\partial]$ and $\sigma^{-n}(\text{lc}_\partial(PL)) = \frac{w}{vp} \text{lc}_\partial(L)$. We then call P a *p-removing operator* for L , and PL the corresponding *p-removed operator*. p is simply called *removable* from L if it is removable at order n for some $n \in \mathbb{N}$.

Example 3.

- (1) In the example from the introduction, we have $L = x(1 - x)D - 1 \in K[x][D]$. An x -removing operator is $P = \frac{1}{x}D$: we have $PL = (1 - x)D^2 - 2D$. Because of $\text{deg}_\partial(P) = 1$ we say that x is removable at order 1.

If P is a p -removing operator then so is QP , for every $Q \in K[x][\partial]$ with

$$\text{gcd}(\text{lc}_\partial(Q), \sigma^{\text{deg}_\partial(P)+\text{deg}_\partial(Q)}(p)) = 1.$$

In particular, note that the definition permits to introduce some new factors w into the leading coefficient while p is being removed. For instance, in our example also $\frac{2-3x}{x}D$ is an x -removing operator for L .

- (2) The definition does not imply that the leading coefficient of a p -removed operator is coprime with $\sigma^n(p)$. In general, it only requires that the multiplicity is reduced. As an example, consider the operator

$$L = x^2(x - 2)(x - 1)D^2 + 2x(x^2 - 3x + 1)D - 2 \in K[x][D]$$

and $p = x$. The operator $P = \frac{x^4 - x^3 - 4x^2 + 2x - 2}{(x - 2)x}D - (x^2 + 5x + 3) \in K(x)[D]$ is a p -removing operator because the leading coefficient of

$$\begin{aligned} PL &= x(x - 1)(x^4 - x^3 - 4x^2 + 2x - 2)D^3 \\ &\quad - (x^6 - 4x^5 - x^4 + 22x^3 - 18x^2 + 18x - 6)D^2 \\ &\quad - 2(x^5 - x^4 - 8x^3 + 8x^2 - 3x + 6)D \\ &\quad + 2(x^2 + 5x + 3) \end{aligned}$$

contains only one copy of p while there are two of them in L . This is called *partial desingularization*. Observe that the definition permits to remove some factors v from the leading coefficient in addition to p .

- (3) In the shift case, or more generally, in an Ore algebra where σ is not the identity, the leading coefficient changes when an operator is multiplied by a power of ∂ from the left. The application of σ^{-n} in the definition compensates for this change. As an example, consider the operator

$$\begin{aligned} L &= x(x + 1)(5x - 2)S^2 - 2x(5x^2 - 2x - 9)S \\ &\quad + (x - 4)(x + 2)(5x + 3) \in K[x][S] \end{aligned}$$

and $p = x + 1$. The operator $P = \frac{5x^3 + 13x^2 - 18x - 24}{(x + 2)(5x + 3)}S - \frac{2(5x^3 + 28x^2 + 23x - 24)}{(x + 2)(5x + 3)}$ is a p -removing operator because the leading coefficient of

$$\begin{aligned}
 PL &= (x + 1)(5x^3 + 13x^2 - 18x - 24)S^3 \\
 &\quad - 2(x + 1)(10x^3 + 21x^2 - 58x + 24)S^2 \\
 &\quad + (25x^4 + 60x^3 - 217x^2 - 84x + 288)S \\
 &\quad - 2(x - 4)(5x^3 + 28x^2 + 23x - 24)
 \end{aligned}$$

does not contain $\sigma(p) = x + 2$. It is irrelevant that it contains $x + 1$.

As indicated in the examples, when removing a factor p from an operator L , Definition 2 allows that we introduce other factors w , coprime to p . We are also always allowed to remove additional factors v besides p . The freedom for having v and w is convenient but not really necessary. In fact, whenever there exists an operator $P \in K(x)[\partial]$ of order n such that $\sigma^{-n}(\text{lc}_\partial(PL)) = \frac{w}{vp} \text{lc}_\partial(L)$, then there also exists an operator $Q \in K(x)[\partial]$ of order n such that $\sigma^{-n}(\text{lc}_\partial(QL)) = \frac{1}{p} \text{lc}_\partial(L)$. To see this, note that by the extended Euclidean algorithm there exist $s, t \in K[x]$ such that $sw + tp = 1$. Set $Q = \sigma^n(sv)P + \sigma^n(t)\partial^n$. Then

$$\begin{aligned}
 \sigma^{-n}(\text{lc}_\partial(QL)) &= sv \sigma^{-n}(\text{lc}_\partial(PL)) + t \sigma^{-n}(\text{lc}_\partial(\partial^n L)) \\
 &= sv \frac{w}{vp} \text{lc}_\partial(L) + \frac{tp}{p} \text{lc}_\partial(L) = \frac{1}{p} \text{lc}_\partial(L),
 \end{aligned}$$

as desired. This argument is borrowed from Abramov et al. (2006). The same argument can also be used to show the existence of operators that remove all the removable factors in one stroke:

Lemma 4. Let $L \in K[x][\partial]$, let $n \in \mathbb{N}$, and let $\text{lc}_\partial(L) = p_1^{e_1} p_2^{e_2} \cdots p_m^{e_m}$ be a factorization of the leading coefficient into irreducible polynomials. For each $i = 1, \dots, m$, let $k_i \leq e_i$ be maximal such that p_i is removable from L at order n . Then there exists an operator $P \in K(x)[\partial]$ of order n such that $\sigma^{-n}(\text{lc}_\partial(PL)) = \frac{1}{p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}} \text{lc}_\partial(L)$.

Proof. By the remark preceding the lemma, we may assume that for every i there exists an operator $P_i \in K(x)[\partial]$ of order n with $P_i L \in K[x][\partial]$ and $\sigma^{-n}(\text{lc}_\partial(P_i L)) = p_i^{-k_i} \text{lc}_\partial(L)$ (i.e., $w = v = 1$).

Next, observe that when p and q are two coprime factors of $\text{lc}_\partial(L)$ which both are removable at order n , then also their product pq is removable at order n . Indeed, if $P, Q \in K(x)[\partial]$ are such that $\deg_\partial(P) = \deg_\partial(Q) = n$, $PL, QL \in K[x][\partial]$, $\sigma^{-n}(\text{lc}_\partial(PL)) = \frac{1}{p} \text{lc}_\partial(L)$, and $\sigma^{-n}(\text{lc}_\partial(QL)) = \frac{1}{q} \text{lc}_\partial(L)$, and if $s, t \in K[x]$ are such that $sq + tp = 1$, then for $R := \sigma^n(s)P + \sigma^n(t)Q$ we have $\sigma^{-n}(\text{lc}_\partial(RL)) = \frac{1}{pq} \text{lc}_\partial(L)$, as desired.

The claim of the lemma now follows by induction on m , taking $p = p_1^{e_1} \cdots p_{m-1}^{e_{m-1}}$ and $q = p_m^{e_m}$. □

4. Desingularization by taking least common left multiples

As outlined in Section 2, the classical algorithm for desingularizing differential operators relies on taking the lclm of the operator to be desingularized with a suitably chosen auxiliary operator. Our contribution consists in a three-fold generalization of this approach: first, we show that it works in every Ore algebra and not just for differential operators, second, we show that almost every operator qualifies as an auxiliary operator in the lclm and not just the particular operator used traditionally, and third, we show that the approach also covers partial desingularization. From the second fact it follows directly that taking the lclm with a random operator of appropriate order removes, with high probability, all the removable singularities of the operator under consideration and not just a given one.

Consider an operator $L \in K[x][\partial]$ in an arbitrary Ore algebra, and let $p \mid \text{lc}_\partial(L)$ be a factor of its leading coefficient. Assume that this factor is removable at order n . Our goal is to show that for almost all operators $A \in K[\partial]$ of order n with constant coefficients the operator $\text{lclm}(L, A)$ is p -removed.

One way of computing the least common left multiple of two operators $L, A \in K[x][\partial]$ with $\deg_\partial(L) = r$ and $\deg_\partial(A) = n$ (and not necessarily with constant coefficients) is as follows. Make

an ansatz with undetermined coefficients $u_0, \dots, u_n, v_0, \dots, v_r$ and compare coefficients of ∂^i ($i = 0, \dots, n+r$) in the equation

$$(u_0 + \dots + u_{n-1}\partial^{n-1} + u_n\partial^n)L = (v_0 + \dots + v_{r-1}\partial^{r-1} + v_r\partial^r)A.$$

This leads to a system of homogeneous linear equations over $K(x)$ for the undetermined coefficients, which has more variables than equations and therefore must have a nontrivial solution. For each solution, the operator on either side of the equation is a common left multiple of L and A .

For most choices of A the solution space will have dimension $\#vars - \#eqns = 1$, and in this case, for every nontrivial solution we have $u_n \neq 0$. In particular the least common left multiple $M = \text{lcm}(L, A)$ has then order $r+n$. The singularities of M are then the roots of $\sigma^n(\text{lc}_\partial(L))$ plus the roots of u_n minus the common roots of u_0, \dots, u_n , which are divided out because we defined the notation $\text{lcm}(L, A)$ to refer to a least common left multiple with polynomial coefficients that share no common factors. It is not obvious at this point why removable factors should appear among these common factors of u_0, \dots, u_n . To see that they systematically do, consider a p -removing operator $P \in K(x)[\partial]$ of order n , and observe that the operators $1, \partial, \dots, \partial^{n-1}, \partial^n$ generate the same $K(x)$ -vector space as $1, \partial, \dots, \partial^{n-1}, P$. If we use the latter basis in the ansatz for the lcm, i.e., do coefficient comparison in

$$(u_0 + \dots + u_{n-1}\partial^{n-1} + u_nP)L = (v_0 + \dots + v_{r-1}\partial^{r-1} + v_r\partial^r)A,$$

then every nontrivial solution vector $(u_0, \dots, u_n, v_0, \dots, v_r)$ of the resulting linear system gives rise to a common left multiple of L and A in $K[x][\partial]$ whose singularities are the roots of $\text{lc}_\partial(PL) = \sigma^n(\frac{1}{p}\text{lc}_\partial(L))$ plus the roots of u_n minus the common roots of u_0, \dots, u_n . This argument shows that the removable factor p will have disappeared in the lcm unless it is reintroduced by u_n . The main technical difficulty to be addressed in the following is to show that this can happen only for very special choices of A . For the proof of this result we need the following lemma.

Lemma 5. *Let F be a field. Let $n, m \in \mathbb{N}$, let $v_1, \dots, v_n \in F^{n+m}$ be linearly independent over F , and let $w_1, \dots, w_m \in F[x_1, \dots, x_n]^{n+m}$ be defined by*

$$w_1 = \begin{pmatrix} x_1 \\ \vdots \\ x_n \\ 1 \\ 0 \\ \vdots \\ \vdots \\ 0 \end{pmatrix}, \quad w_2 = \begin{pmatrix} 0 \\ x_1 \\ \vdots \\ x_n \\ 1 \\ 0 \\ \vdots \\ \vdots \\ 0 \end{pmatrix}, \quad \dots, \quad w_m = \begin{pmatrix} 0 \\ \vdots \\ \vdots \\ \vdots \\ 0 \\ x_1 \\ \vdots \\ \vdots \\ x_n \\ 1 \end{pmatrix}.$$

Then $\Delta := \det(w_1, \dots, w_m, v_1, \dots, v_n) \in F[x_1, \dots, x_n]$ is not the zero polynomial.

Proof. Simultaneous induction on n and m : We show that the lemma holds for (n, m) if it holds for $(n-1, m)$ and for $(n, m-1)$.

As induction base, observe first that the lemma holds for $n=1, m$ arbitrary: proceeding from the bottom up, use the columns w_1, \dots, w_m to eliminate the nonzero entries of v_1 , of which there must be at least one. Each elimination of some coordinate of v_1 introduces a multiple of x_1 in the next coordinate. Since each coordinate except for the first can be reduced by some w_i , this procedure turns v into a vector of the form $(p, 0, \dots, 0)^T$, for some nonzero polynomial $p \in F[x_1]$. We thus have $\Delta = \pm p \neq 0$.

Observe secondly, still for the induction base, that the lemma also holds for n arbitrary and $m=1$. To see this, note that the coordinates $1, x_1, \dots, x_n$ of w_1 are linearly independent over F . By Laplace-expanding the determinant Δ along w_1 , we see that it is zero if and only if all the $n \times n$ -minors of $(v_1, \dots, v_n) \in F^{(n+1) \times n}$ are zero. But in this case, by Cramer's rule, v_1, \dots, v_n would be linearly dependent, which by assumption they are not. So $\Delta \neq 0$.

Now let $(n, m) \in \mathbb{N}^2$ with $n \geq 2, m \geq 2$ be given. Let $v_1, \dots, v_n \in F^{n+m}$ be linearly independent. Write $v_i = (v_{1,i}, \dots, v_{n+m,i})$ for the coefficients. For a vector $u \in F^{n+m}$, we write \bar{u} for the vector in F^{n+m-1} obtained from u by chopping off the first coordinate.

Case 1. $v_{1,1} = v_{1,2} = \dots = v_{1,n} = 0$. In this case, the vectors $\bar{v}_i \in F^{n+(m-1)}$ must be linearly independent. By expanding along the first row, we have

$$\Delta = x_1 \det(\bar{w}_2, \dots, \bar{w}_m, \bar{v}_1, \dots, \bar{v}_n).$$

The determinant on the right is nonzero by applying the lemma with n and $m - 1$. Therefore the determinant on the left is also nonzero.

Case 2. If at least one of the $v_{1,j}$ is nonzero, then we may assume without loss of generality that $v_{1,1} = 1$ and $v_{1,2} = v_{1,3} = \dots = v_{1,n} = 0$, by performing suitable column operations on $(v_1, \dots, v_n) \in F^{(n+m) \times n}$. Then the vectors $\bar{v}_2, \dots, \bar{v}_n \in F^{(n-1)+m}$ obtained from the v_i by chopping the first coordinate are linearly independent. Expanding along the first row, we now have

$$\Delta = x_1 \left[\text{[poly]} \right] + v_{1,1} \times \begin{vmatrix} x_2 & x_1 & 0 & \cdots & 0 & v_{2,2} & \cdots & v_{2,n} \\ x_3 & x_2 & \ddots & \ddots & \vdots & \vdots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 & \vdots & & \vdots \\ \vdots & & \ddots & \ddots & x_1 & \vdots & & \vdots \\ x_n & & & \ddots & x_2 & \vdots & & \vdots \\ 1 & \ddots & & & x_3 & \vdots & & \vdots \\ 0 & \ddots & \ddots & & \vdots & \vdots & & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots & \vdots & & \vdots \\ \vdots & & \ddots & \ddots & x_n & \vdots & & \vdots \\ 0 & \cdots & \cdots & 0 & 1 & v_{n+m,2} & \cdots & v_{n+m,n} \end{vmatrix}.$$

By setting $x_1 = 0$, the first term on the right hand side disappears, and so do the entries x_1 in the determinant of the second term. By applying the lemma with m and $n - 1$, the determinant on the right with x_1 set to zero is a nonzero polynomial in x_2, \dots, x_n . Since also $v_{1,1} \neq 0$, the whole right hand side is nonzero for $x_1 = 0$. Consequently, when x_1 is not set to zero, it cannot be the zero polynomial. □

Theorem 6 (Main result). Let $K[x][\partial]$ be an Ore algebra, let $L \in K[x][\partial]$ be an operator of order r , and let $n \in \mathbb{N}$. Let $p \in K[x]$ be an irreducible polynomial which appears with multiplicity e in $\text{lc}_\partial(L)$ and let $k \leq e$ be maximal such that p^k is removable from L at order n . Let $A = a_0 + a_1\partial + \dots + a_{n-1}\partial^{n-1} + \partial^n$ in $K[a_0, \dots, a_{n-1}][\partial]$, where a_0, \dots, a_{n-1} are new constants, algebraically independent over K . Then the multiplicity of $\sigma^n(p)$ in $\text{lc}_\partial(\text{lclm}(L, A))$ is $e - k$.

Proof. Let $P_0, \dots, P_n \in K(x)[\partial]$ be such that each P_i has order i and removes from L all the factors of $\text{lc}_\partial(L)$ that can possibly be removed by an operator of order i . Such operators exist by Lemma 4. Consider an ansatz

$$u_0 P_0 L + u_1 P_1 L + \dots + u_n P_n L = v_0 A + v_1 \partial A + \dots + v_r \partial^r A$$

with unknown $u_i, v_j \in K[a_0, \dots, a_{n-1}][x]$. Compare coefficients with respect to powers of ∂ on both sides and solve the resulting linear system using Cramer's rule. This gives a polynomial solution vector with

$$u_n = \det([P_0 L], [P_1 L], \dots, [P_{n-1} L], [A], [\partial A], \dots, [\partial^{r-1} A]),$$

where the notation $[U]$ refers to the coefficient vector of the operator U (padded with zeros, if necessary, to dimension $r + n$).

If $\sigma^n(p) \mid u_n$, then the columns of the determinant are linearly dependent when viewed as elements of $F[a_0, \dots, a_{n-1}]$ with $F = K[x]/\langle \sigma^n(p) \rangle$. Then Lemma 5 implies that already $[P_0L], \dots, [P_{n-1}L]$ are linearly dependent modulo $\sigma^n(p)$. In other words, there are polynomials $u_0, \dots, u_{n-1} \in K[x]$ of degree $< \deg(p)$, not all zero, such that the linear combination $u_0P_0L + \dots + u_{n-1}P_{n-1}L$ has content $\sigma^n(p)$. If $d \in \{0, \dots, n-1\}$ is maximal such that $u_d \neq 0$, then this means that $\frac{1}{\sigma^n(p)}(u_0P_0 + \dots + u_dP_d)$ is an operator of order d which removes from L one factor $\sigma^{n-d}(p)$ more than P_d does, in contradiction to the assumption that P_d removes as much as possible.

This proves that $\sigma^n(p) \nmid u_n$, and in particular $u_n \neq 0$. Since P_n is assumed to remove all removable factors, and in particular k copies of p , and since u_n does not re-introduce any copy of p , it follows that the multiplicity of $\sigma^n(p)$ in $\text{lc}_\partial(u_nP_nL) = \text{lc}(\text{lcm}(L, A))$ is $e - k$, as claimed. \square

The theorem continues to hold when the indeterminates a_0, \dots, a_{n-1} are replaced by values in K which do not form a point on the zero set of the determinant polynomial $u_n \bmod \sigma^n(p)$, as discussed in the proof. As this is not the zero polynomial and we assume throughout that K has characteristic zero, it follows that almost all choices of $A \in K[\partial]$ will successfully remove all the factors of $\text{lc}_\partial(L)$ that are removable at order $\deg_\partial(A)$.

The theorem thus justifies the following very simple probabilistic algorithm for removing, with high probability, as many factors as possible from a given operator $L \in K[x][\partial]$ at a given order n :

- Pick an operator $A \in K[\partial]$ of order n at random.
- Return $\text{lcm}(L, A)$.

This is a Monte Carlo algorithm: it always terminates but with low probability may return an incorrect answer. For a deterministic algorithm, don't take the operators A at random but use an operator with symbolic constant coefficients a_0, \dots, a_{n-1} , as in the theorem. The leading coefficient of lcm will then have all removable factors removed, and some additional factors involving the symbolic coefficients. Now instantiate these coefficients with some elements of K for which they don't evaluate to one of the factors of $\sigma^n(\text{lc}_\partial(L))$. Almost any choice will do. By taking the choices from some enumeration of \mathbb{Z}^n , for example, it is guaranteed that we will encounter a choice that works after finitely many attempts.

The Monte Carlo version of the algorithm is included in the new `ore_algebra` package for Sage (Kauers et al., 2015), and works very efficiently thanks to the efficient implementation of least common left multiples also available in this package. This package has been used for the calculations in the following concluding examples. The computation time for all these examples is negligible.

Example 7.

(1) For $L \in \mathbb{Q}[x][D]$ from Example 1 and the “randomly chosen” operator $A = D^2 + D + 1$ we have

$$\begin{aligned} \text{lcm}(L, A) &= (x^7 - 4x^6 + 6x^5 - 4x^4 + x^3 + 6x - 6)D^4 \\ &\quad - (2x^6 - 9x^5 + 15x^4 - 11x^3 + 3x^2 - 24)D^3 \\ &\quad - (x^7 - 4x^6 + 6x^5 - 4x^4 + x^3 + 6x - 6)D \\ &\quad + (2x^6 - 9x^5 + 15x^4 - 11x^3 + 3x^2 - 24). \end{aligned}$$

This is not the same result as in Example 1, but it does have the required property $x \nmid \text{lc}_\partial(\text{lcm}(L, A))$.

(2) This is an example for the recurrence case. Let

$$\begin{aligned} L &= 2(x+3)^2(59x+94)S^3 - (2301x^3 + 15171x^2 + 32696x + 22876)S^2 \\ &\quad - 5(59x^3 + 330x^2 + 600x + 359)S - (59x + 153)(x+1)^2. \end{aligned}$$

Among the factors of $(x + 3)$ and $(59x + 94)$ of the leading coefficient, the latter is removable at order 1 and the former is not removable. Accordingly, for the “randomly chosen” operator $A = S - 2$ we have

$$\text{lclm}(L, A) = 2(x + 4)^2(8909x^3 + 57087x^2 + 119629x + 81711)S^4 + (\dots)S^3 + (\dots)S^2 + (\dots)S + (\dots),$$

where (\dots) stands for some other polynomials. Note that the leading coefficient is coprime to $\sigma(59x + 94) = 59x + 153$.

- (3) As an example for partial desingularization, consider the operator $L = x^3D^3 - 3x^2D^2 - 2xD + 10 \in \mathbb{Q}[x][D]$. Of the three copies of x in the leading coefficient, one is removable at order 2, another one at order 4, and the third is not removable. In perfect accordance, we find for example

$$\begin{aligned} \text{lc}_\partial(\text{lclm}(L, D + 2)) &= x^3(4x^3 + 6x^2 - 2x - 5), \\ \text{lc}_\partial(\text{lclm}(L, D^2 + 1)) &= x^2(x^6 + 10x^4 + 40x^2 + 80), \\ \text{lc}_\partial(\text{lclm}(L, D^3 + 3D^2 - 1)) &= x^2(x^8 - 30x^6 + \dots + 2160x + 1920), \\ \text{lc}_\partial(\text{lclm}(L, D^4 - D^2 + 1)) &= x(x^{10} - 10x^8 + 120x^6 - 720x^4 - 3200), \\ \text{lc}_\partial(\text{lclm}(L, D^5 + D - 1)) &= x(x^{12} - 3x^{11} + \dots + 25600x - 22400). \end{aligned}$$

- (4) There are unlucky choices for A . For example, consider

$$\begin{aligned} L &= (x - 7)(x^2 - 2x - 12)S^2 - (3x^3 - 23x^2 - 23x + 291)S \\ &\quad + 2(x - 6)(x^2 - 13) \in \mathbb{Q}[x][S]. \end{aligned}$$

The factor $x - 7$ is removable, as can be seen, for example, from the fact that $\text{lc}_\partial(\text{lclm}(L, S - 1)) = 2x^2 - x - 51$ is coprime to $\sigma(x - 7) = x - 6$. However, if we take $A = S - \frac{9}{4}$, then

$$\begin{aligned} \text{lclm}(L, A) &= 4(x - 7)(x - 6)(5x - 28)S^3 \\ &\quad - (x - 7)(3092 - 1138x + 105x^2)S^2 \\ &\quad + (x - 5)(6081 - 2080x + 175x^2)S \\ &\quad - 18(x - 6)(x - 5)(5x - 23), \end{aligned}$$

which has $x - 6$ in the leading coefficient. (It is irrelevant that also $x - 7$ appears as a factor.)

- (5) Finally, as an example in an unusual Ore algebra, consider $\mathbb{Q}[x][\partial]$ with $\sigma: \mathbb{Q}[x] \rightarrow \mathbb{Q}[x]$ defined by $\sigma(x) = x^2$ and $\delta: \mathbb{Q}[x] \rightarrow \mathbb{Q}[x]$ defined by $\delta(x) = 1 - x$. Let

$$L = (2x + 1)\partial^2 + (x^2 + 3x - 1)\partial - (2x^4 + 2x^3 + x^2 + 1).$$

The factor $2x + 1$ is removable at order 1. For example, for $A = \partial - 1$ we find that $\text{lclm}(L, A)$ equals

$$\begin{aligned} &(2x^3 + 4x^2 + 4x - 1)\partial^3 - (2x^6 - x^4 - 4x^3 - 3x^2 + x + 5)\partial^2 \\ &\quad - (2x^9 + 4x^8 + 6x^7 + 4x^6 + 2x^5 + 3x^4 + 2x^3 + 3x^2 + 3x - 2)\partial \\ &\quad + (2x^9 + 4x^8 + 6x^7 + 6x^6 + 2x^5 + 2x^4 - 4x^3 - 4x^2 + 4). \end{aligned}$$

As expected, the leading coefficient does not contain $\sigma(\text{lc}_\partial(L)) = 2x^2 + 1$.

References

- Abramov, Sergei A., Barkatou, Moulay A., van Hoeij, Mark, 2006. Apparent singularities of linear difference equations with polynomial coefficients. *Appl. Algebra Eng. Commun. Comput.* 17, 117–133.
- Abramov, Sergei A., Le, Ha Q., Li, Ziming, 2005. Univariate Ore polynomial rings in computer algebra. *J. Math. Sci.* 131, 5885–5903.
- Abramov, Sergei A., van Hoeij, Mark, 1999. Desingularization of linear difference operators with polynomial coefficients. In: *Proceedings of ISSAC'99*, pp. 269–275.
- Abramov, Sergei A., van Hoeij, Mark, 2003. Sets of poles of solutions of linear difference equations with polynomial coefficients. *Comput. Math. Math. Phys.* 43 (1), 57–62.
- Bostan, Alin, Chyzak, Frédéric, Li, Ziming, Salvy, Bruno, 2012. Fast computation of common left multiples of linear ordinary differential operators. In: *Proceedings of ISSAC'12*, pp. 99–106.
- Bronstein, Manuel, Petkovšek, Marko, 1996. An introduction to pseudo-linear algebra. *Theor. Comput. Sci.* 157 (1), 3–33.
- Chen, Shaoshi, Kauers, Manuel, 2012a. Order-degree curves for hypergeometric creative telescoping. In: *Proceedings of ISSAC'12*, pp. 122–129.
- Chen, Shaoshi, Kauers, Manuel, 2012b. Trading order for degree in creative telescoping. *J. Symb. Comput.* 47 (8), 968–995.
- Chen, Shaoshi, Kauers, Manuel, Jaroschek, Maximilian, Singer, Michael, 2013. Desingularization explains order-degree curves for Ore operators. In: *Proceedings of ISSAC'13*, pp. 157–164.
- Chyzak, Frédéric, Dumas, Philippe, Le, Ha, Martin, Jose, Mishna, Marni, Salvy, Bruno, in preparation. Taming apparent singularities via Ore closure.
- Ince, E.L., 1926. *Ordinary Differential Equations*. Dover.
- Kauers, Manuel, Jaroschek, Maximilian, Johansson, Fredrik, 2015. Ore polynomials in Sage. In: *Computer Algebra and Polynomials*. In: *Lect. Notes Comput. Sci.*, vol. 8942, pp. 105–125.
- Ore, Øystein, 1933. Theory of non-commutative polynomials. *Ann. Math.* 34 (3), 480–508.
- Schlesinger, Ludwig, 1895. *Handbuch der Theorie der linearen Differentialgleichungen*. Teubner.